



Early and Middle Pleistocene Faunal and Hominins Dispersals through Southwestern Asia

Citation

Bar-Yosef, Ofer and Miriam Belmaker. Forthcoming. Early and Middle Pleistocene faunal and hominins dispersals through Southwestern Asia. Quaternary Science Reviews 29.

Published Version

doi:10.1016/j.quascirev.2010.02.016

Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:4270472>

Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Open Access Policy Articles, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#OAP>

Share Your Story

The Harvard community has made this article openly available.
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)

**Early and Middle Pleistocene Faunal and Hominins Dispersals through
Southwestern Asia**

Ofer Bar-Yosef and Miriam Belmaker

Department of Anthropology

Harvard University

11 Divinity Avenue

Cambridge MA 02138

Phone ++ 1 617 495 1279

Fax ++ 1 617 496 8041

Abstract

This review summarizes the paleoecology of the Early and Middle Pleistocene of southwestern Asia, based on both flora and fauna, retrieved from a series of ‘windows’ provided by the excavated sites. The incomplete chrono-stratigraphy of this vast region does not allow to accept the direct chronological correlation between the available sites and events of faunal and hominin dispersals from Africa. It also demonstrates that hominins survived in a mixed landscape of open parkland with forested surrounding hills. In addition, the prevailing environmental conditions are not sufficient to explain the differences between ‘core and flake’ and the Acheulian industries that probably reflect the learned traditions of different groups of hominins successful adaptations to new ecological niches away from the African savanna. The current distribution of lithic industries across Eurasia is undoubtedly incomplete due to lack of cultural continuities as well as paucity of field research in several sub-regions. This observation supports the contention that what we view as a constant stream of migrants was actually interrupted many times. The continuous occupation of southwestern Asia by the makers of the Acheulian is in contrast with neighboring regions such as the Iranian plateau and Eastern Europe. A more complex model is required to explain the Eurasian archaeological-cultural mosaic recorded in Eurasia.

Introduction

The effect of climate change on the tempo and mode of early hominids dispersals from Africa during the Early and Middle Pleistocene is one of the main interests in paleoanthropology and Paleolithic archaeology (Behrensmeyer, 2006). It has been suggested that the expansion of savanna environments during the Early Pleistocene allowed for the first ‘Out of Africa I’ dispersal (Dennell, 2004; Martínez-Navarro, 2004) and that humans dispersed into Western Europe during warmer periods and that the dispersal was highly influenced by climate rather than culture, which played only a minor role (Agustí et al., 2009a). However, these theories assume as a working hypotheses that the presence of hominins and faunal present in the paleontological and archaeological record is continuous. Here we would like to focus on southwestern Asia that served as the main corridor for hominin dispersal during the Early and Middle Pleistocene and address the assertion that we can test hominids co-dispersed with other taxa and the extent of the response to climatic forcing. We will propose that there is no correlation between hominid and faunal dispersals and suggest that early *Homo* dispersals were not directly controlled by climatic forcing.

Geological and archaeological investigations in southwestern Asia produced a few “windows” into the geological sequence of the Early and Middle Pleistocene. Most of these “windows” are known from excavations of archaeological sites. Hence, in order to provide a continuous narrative of hominin and faunal dispersals we need to combine the chronologically patchy information with the physiographic variability of this vast region, as well as the incomplete paleo-climatic conditions that determined the distribution of resources in each sub-region.

The sub-regions of southwestern Asia are commonly identified from north to south and west to east as follows: Anatolia, the southern Caucasus, The Zagros mountains and the Iranian plateau, the Levant and the Arabian peninsula. Among these, due to many years of field and laboratory research, the better known is the Levant. It also has the advantage that researchers from various schools and institutions conducted surveys of Quaternary deposits, drilling boreholes for water and pollen, and archaeological excavations of Lower Paleolithic occupations. Thus the accumulated information creates an imbalance between the areas, some of which like the Iranian and the Anatolian plateaus are of larger area than the Levant but yet poorly known. We are therefore forced to base our review on the Levantine sequence and touch upon, in a more cursory fashion, the other sub-regions.

The Levant is a unique biogeographic entity within southwestern Asia. It lies at the crossroads of Africa and Eurasia and has more lush environments compared to the alternative dispersal corridor on the southern fringes of the Arabian Peninsula. This land bridge emerged during the Miocene as a continuous terrestrial belt that allowed various plants and animal taxa to migrate through in either direction (Thomas, 1985; Tchernov, 1988; Bar-Yosef, 1994,1998a). The southern path through the margins of the Arabian Peninsula could be crossed at the Bab el-Mandeb straits during low sea level stands, and along the area that receives the summer Indian Ocean monsoon to be followed by a passageway through the Hormuz straits into the southern coast of Iran (Fig. 1), and a northern path via the Nile river and the Sinai Peninsula.

During the Middle Pleistocene both land and sea bridges were potential dispersal routes for hominins (Derricourt, 2006). However, the available archaeological evidence

was retrieved only from the Levant with a few indications from the southern route (Amirkhanov, 1991). During the Late Pliocene and Early Pleistocene there is no evidence for a land bridge over the Bab el Mandeb straits. Indeed, during this time period, dispersals of most animal and plants seem to have been limited to the terrestrial Northern corridor of the Levant.

It is well-known that dispersals of early *Homo ergaster/etectus* from Africa across Eurasia during the Early and Middle Pleistocene are interpolated from isolated archaeological and fossil sites (e.g., Gabunia and Vekua, 1995; Larick and Ciochon, 1996; Bar-Yosef, 1998a; Potts, 1998; Arribas and Palmqvist, 1999; Bar-Yosef and Belfer-Cohen, 2001; Potts, 2002; Antón and Swisher, 2004; Martínez-Navarro, 2004; Lordkipanidze et al., 2007). The reasons for these long-range migrations, whether in a form of “leap frog” expansions or gradual diffusion, are formulated as plausible hypotheses, open to alternative interpretations. Moreover, the advantages for leaving Africa are not easily testable (e.g., Bar-Yosef and Belfer-Cohen, 2001). One may use the biogeographic ranges outside Africa, from the Iberian peninsula to East and Southeast Asia, occupied by early hominins, to calculate the rate of movements and the chance for long term survival of early mobile hunter-gatherers. Currently it seems that it took ca. 0.5 Ma to reach Western Europe by ca. 1.3 Ma (Agustí et al., 2000) and much less time to get to ca. 1.66-1.7 Ma to East and Southeast Asia (Zhu et al., 2003; Antón and Swisher, 2004). One of the major questions is how African species adapted to the ecological variability in Eurasia and the role the ‘culture’, as recorded in the stone tools, assisted (or not) in the survival of lineages in the new territories.

For the purpose of making the most parsimonious explanations about hominin adaptations, and indicating where and when continuity is interrupted, probably by the demise of some groups, we provide a brief review of current climatic and vegetation conditions, a survey of the main Lower Paleolithic sites, followed by a summary of the vegetation and faunal records as a basis for environmental reconstruction and long distance correlations with known climatic events, and evaluation of hominin dispersals through this region.

Current Climatic Conditions

The dominant physiographic feature of southwestern Asia is the combination of mountains, plateaus, alluvial plains and desert landscapes including oases. The Anatolia plateau is surrounded by mountains ranges in the north and the south where the Taurus Mountains arching eastward into the Zagros Mountains. The Levant, along the eastern Mediterranean coast has several parallel features. The narrow coastal plain in the north widens in the south, the mountain ranges create the main watershed of this sub-region, the Anuq-Beqa'a-Jordan Rift Valley stretches from north to south, and the Trans-Jordanian or Syro-Arabia desert plateau descends eastward and southward. The Euphrates and the Tigris rivers and their tributaries cross the Mesopotamian plain, which is bordered by the Zagros Mountains and the Iranian plateau. The latter was and is mainly desert strewn by oases.

Today's climate of the region is dominated by cold winters with the precipitation brought by the westerlies. Snow falls only in the higher mountains, and in particular in eastern Turkey and the northwestern Zagros. Summers are hot and dry. There is a wide

range of variation across the region. Thus, winter temperatures are milder in the coastal belts and more severe inland or at higher elevations and precipitation is affected by distance from the Mediterranean Sea and the local sub-regional altitude.

The East Mediterranean region is located between the more temperate European climatic zone in the north and the hyper arid regions of the Saharo-Arabian desert belt in the south. Since the synoptic hydrological conditions may affect the entire southwestern Asia, the question is whether and how warm to cold and humid to dry oscillations affected the Levant since the Early Pleistocene (Frumkin and Stein, 2004).

Models of Pleistocene climates attempted to explain the paleoclimatic record of the Levant. Earlier Butzer (1958) suggested a high correlation between the local intensification of the westerly circulation of the Mediterranean during glacial periods and the deep Cyprus Lows that would explain the increase in rainfall during glacial times. Similar views were expressed by others (e.g., Horowitz, 1979). Recently, the extent the increase of annual precipitation during North Hemispheric glaciations affected the Levant, and whether it extended into the currently dry desert belt in the south has been a topic of debate (Enzel et al., 2008 and references therein).

The climatic conditions and physiographic conditions determined the distribution of the phytogeographic belts (Zohary, 1973; Danin, 1988), and the most frequently encountered mammals in each region (e.g., Harrison and Bates, 1991; Harrison, 1972):

1. The Pontic province – the ‘Euro-Siberian’ region corresponds to the Palearctic deciduous and coniferous forests. It stretches from the southern edge of the Black Sea in northern Turkey as well as relict enclaves in the south in the Anti-Taurus Mountains, the Syrian Amanus range and northern Lebanon. Part of the Pontic includes the Hyrcanian

province of the coastal mountains of the southern Caspian Sea that descends into Iraqi-Kurdistan and northern Syria. It included deciduous species of maple (*Acer*), birch (*Bitula*), hazelnut (*Corylus*), beech (*Fagus*), ash (*Fraxinus*), lime (*Tilia*) and elm (*Ulmus*) that have a clear Arcto-Tertiary origin. Other evergreen species that occur in southern Europe are present such as the boxwood (*Buxus sempervirens*), holly (*Ilex aquifolium*) and yew (*Taxus baccata*).

2. Irano-Turanian province – covers the sub-regions of the Anatolian plateau, the Syrian Desert, the Iranian plateau, and stretches into central Asia and the Gobi Desert. The climate here is dominated by extremely cold winters and very hot summers with little to no rain fall in the winter. It is dominated by dwarf scrubland and steppes vegetation.

3. The Saharo-Arabian province stretches south of the Mediterranean basin and extends from Mauritania to the Red Sea incorporating the Sahara desert. This vegetation is open xeromorph dwarf scrubland and desert plant associations.

4. The Mediterranean province spreads around the Mediterranean Sea and includes a combination of *maquis* and forests dominated by oak (*Quercus*), Olive (*Olea*) and almonds (*Amgdalus communis*).

5. The Sudano-Zambesian province is confined to the Jordan Valley and preserves relicts of typical African subtropical savanna species such as the acacias.

Southwestern Asia was frequented by fauna from three biogeographic provinces: Palaearctic, Oriental and Ethiopian in different proportions. Here we include a description of extant fauna, all species dating to the early Holocene that predates the effects of agriculture, horticulture and hunting. Thus most of the data comes from archaeological excavations.

167 Many taxa are similar over two or more provinces giving the entire Near East a
168 coherent faunal community (Harrison and Bate, 1992). Two main regions can be
169 observed: The first includes the Mediterranean, Pontic and Iranian plateau provinces with
170 Palearctic taxa, and the second includes the mesic Mediterranean, Arabian provinces with
171 Ethiopian elements. Taxa common to all regions include *Hyaena hyaena*, *Felis silvestris*,
172 *Felis chaus*, *Panthera pardus*, *Acinonyx jubatus*, *Lepus capensis*, *Hystrix indica*, *Rattus*
173 *rattus*, *Rattus norvegicus*, *Mus macedonicus*. Palearctic fauna which are found in the
174 Mesic regions of the Mediterranean, the Pontic and much of the Iranian plateau include
175 *Dama mesopotamica*, *Capreolus capreolus*, *Cervus elaphus*, *Sus scrofa*, *Canis aureus*,
176 *Canis lupus*, *Vulpes vulpes*, *Panthera pardus*, *Ursus arctos*, *Martes foina*, *Vormela*
177 *peregrina*, *Mustela nivalis*, *Meles meles*, *Lutra lutra*, *Sciurus anomalus*, *Spalax*
178 *leucodon*, *Apodemus mystacinus*, *Apodemus sylvaticus*, *Cricetulus migratorius*, *Meriones*
179 *tristrami* and *Microtus guentheri*. The Ethiopian taxa unique to the Arabian region are
180 *Vulpes rueppellii*, *Vulpes cana* and *Vulpes zerda*, *Mellivora capensis*, *Genetta felina*,
181 *Felis margarita*, *Caracal caracal*, *Procyon capensis*, *Acomys cahirinus*, *Eliomys*
182 *melanurus*, *Gerbillus nanus*, *Gerbillus dayurus*, *Sekeetamys calurus*, and *Psammomys*
183 *obesus*.

184 Nonetheless, while some genera are common throughout the entire region, the
185 species within each genus differ across the provinces. Thus, the unique species that
186 appear in each province give its unique affinities. For example, the caprine in the
187 Caucasus is *Capra caucasica* while the dominant *Capra* in the Iranian plateau is *Capra*
188 *aegagrus* and in the Mediterranean and Arabia is *Capra ibex*. The gazelle in the semi arid
189 regions of the Mediterranean is *Gazella gazelle*, in Arabia it is *Gazella dorcas* and in the

Iranian plateau is *Gazella subgotarossa*. Furthermore, other species are unique to each region. For example, species unique to the Pontic region are *Lynx Lynx*, *Apodemus flavicollis*, and *Mesocricetus auratus*. Species unique to the Iranian plateau are the *Ovis ammon*, *Allactaga euphratica*, *Dryomys nitedula*, *Calomyscus bailwardi*, *Meriones persicus* and *Ellobius fuscocapllius*. Species unique to Arabia are *Oryx leucoryx*, *Tragelaphus imbris*, *Jaculus jaculus*, *Arvicanthis niloticus* and *Gerbillus gerbillus*.

However, it is most important to remember that the home ranges of the species is fluid rather than static, and are constantly changing dependant on climatic and environmental changes and inter and intra species competition. Therefore, the presence-absence of faunal elements correspond to the vegetation conditions, and disperse when resources diminish during worsening climatic conditions that could also result in extinction (Bennett, 1997). Thus in the following sections, we will discuss the changes in the distribution of botanical and faunal compositions across the Early and Middle Pleistocene in relation to the biogeographic provinces of southwestern Asia.

Lower and Middle Pleistocene sites in Southwestern Asia

The series of sites briefly described below form the isolated windows we have in southwestern Asia into this long period of the Early and Middle Pleistocene. In the absence of radiometric dating and/or paleomagnetic information the ages of several of the localities rely on faunal correlations within the region including long distance comparisons with European sites. Indeed, most of the Pleistocene data was acquired through archaeological excavations while the number of assemblages derived from natural accumulations is limited to a few Pliocene sites in the Caucasus region such as

213 Kvabebi (Vekua and Lordkipadinze, 2008; Agustí et al., 2009b) and Bethlehem in the
214 Levant (see below).

215 There are special localities in the Levant that were thought to provide evidence
216 for early presence of hominins, but a close examination do not stand up to the
217 geochronological criteria. The mammalian fauna of Bethlehem is the oldest Plio-
218 Pleistocene assemblage known from the Levant. This was an animal bone accumulation
219 incorporated in a deposit of flint fragments and clay discovered when a local farmer was
220 digging a well (Gardner and Bate, 1937). The small collection of flints was originally
221 thought to include human artifacts but they have since been shown to be natural (Hooijer,
222 1958; Clark, 1961).

223 Another claim for an Early Pleistocene site named Yiron where core and flakes
224 have been retrieved from the gravels in a major crevice. Unfortunately the Yiron basalt
225 on the plateau did not cover this location and thus the date of the lava flow of 2.4 Ma
226 (Ronen et al., 1980; Ronen, 1991a), is unrelated to the human made lithics.

227 The ‘Erq el Ahmar Formation (Horowitz, 1979) is located about 14 km south of
228 the Sea of Galilee, in the Jordan Valley and accumulated prior to the ‘Ubeidiya Formation.
229 The polarity sequence is correlated with late Gauss and early Matuyuma chron, within
230 which the Olduvai subchron was identified (Ron and Levi, 2001). Unfortunately the
231 artifacts mentioned in the different papers were found on the surface, except for a few
232 retrieved from a fluvial conglomerate of an unknown stratigraphic position (Tchernov,
233 1995). Hence, the attribution of these to the Olduvai subchron is only tentative and
234 cannot serve as a well-established evidence for hominin presence prior to Dmanisi (Bar-
235 Yosef, 1998a).

The subsequent analysis of faunal turnovers in the Levant and the Caucasian region is reported in relation to the cultural definitions, and supplemented by the information from dated archaeological contexts (Table 1). It is followed by information from the more poorly known regions of Anatolia, the Iranian plateau and the Arabian Peninsula; For the sake of clarity, we briefly present the major sites that contain archaeological remains and their cultural attributions in the Caucasian and the Levant regions according to their accepted chronology and regardless of their geographic location (Fig. 1). Numerous prehistoric occurrences were assigned to the Early and Middle Pleistocene that did not yield fossil bones or datable substances or deposits amenable to paleomagnetic readings. On morphological grounds the collected or excavated artifacts were attributed to the Acheulian sequence such as the Early Acheulian occurrences at Nahal Zihor in the Negev (Ginat et al., 2003) or the Late Acheulian in Ma'ayan Baruch in the Upper Jordan Valley. In summarizing the archaeological information we do not describe the faunal and floral remains found in each site in this section, as they will be discussed in length in the following one. The condensed information will allow us to demonstrate below that the assumptions concerning continuous hominin dispersals into Eurasia relies on a series of speculations and that the proposed timing of prehistoric migration is not well tied with the transitions among the faunal assemblages.

Dmanisi

The site of Dmanisi is located on a basaltic block bordered by two tributaries of the larger Kura River. The prehistoric deposits were target for systematic excavations

(Vekua, 1987; Dzaparidze et al., 1989; Gabunia and Vekua, 1990, 1995; Liubin and Bosinski, 1995; Liubin and Bosinski, 1995; Lordkipanidze et al., 2007; Jöris, 2008; Rightmire et al., 2006). The geoarchaeological study concluded that entire sequence at Dmanisi could be dated to the Olduvai subchron and immediately after as shown by the normal polarity of the lower lava flow (1.8 ± 0.1 Ma) and the infillings of the upper deposit that date to 1.77 Ma. Several human skulls and postcranial bones are attributed to *Homo erectus/ergaster* (Gabunia and Vekua, 1995; Rightmire et al., 2006; Lordkipanidze et al., 2007). The lithics of Dmanisi are identified as core and flake industry sometimes called “pre-Oldowan” or Mode 1 (de Lumley et al., 2005; Jöris, 2008). The Dmanisi core and flake industry marks the presence of hominins that did not practice the production of handaxes (bifaces). It indicates that handaxes were not a necessity for the colonization of Eurasia. The Dmanisi faunal assemblage is earlier than the faunas of Sènèze and Le Coupet, and thus also earlier than ‘Ubeidiya (Tchernov, 1987).

‘Ubeidiya

‘Ubeidiya is situated on the edge of the western escarpment of the Jordan Rift Valley and its geological structure was uncovered through a series of excavations (already reported in details) an anticline with several undulations disturbed by a few faults (Bar-Yosef and Tchernov, 1972; Bar-Yosef and Goren-Inbar, 1993). The exposed tilted layers were numbered from those observed as the earliest to the latest over a total thickness of 154m. The sequence was subdivided into four cycles: two mostly limnic (Li and Lu) and two essentially terrestrial (Fi and Fu).

The raw materials used for manufacturing artifacts were lava (basalt), flint, and

limestone. The basalt nodules occurred as pebbles, cobbles, boulders, and scree components; the limestone was available as much more sparse cobbles within the beach and wadi deposits; and the flint is found in the same environments as small pebbles and cobbles. The 'Ubeidiya hominins employed each type of rock to shape a different type of object (Bar-Yosef and Goren-Inbar, 1993). Core-choppers and light-duty tools were made of flint, spheroids mainly of limestone, and the handaxe group from basalt, with a few of flint and limestone. There is a direct correlation between the size of the tool-category and the type of raw material. Although basalt is the most common rock and found in every lithological facies at 'Ubeidiya the most abundant object is the core-chopper, which is made of flint and its detached pieces (flakes). It seems that the lithic assemblages from the lower most layers in the sequence (K/III-12, III-20-22, II-23-24) contain an abundance of core-choppers, polyhedrons and spheroids but lack bifaces, except for one trihedral. The samples are large enough to suggest that they may indicate the presence of an early group of hominins that did not produce bifaces. The overlying assemblages contain bifaces in varying frequencies and can be called Early Acheulian (Bar-Yosef and Goren-Inbar, 1993). As the presence or absence of bifaces is taken to designate different groups of hominins one may speculate that as in Dmanisi, the first hominins to arrive at 'Ubeidiya, were not the bearers of the Acheulian.

Among the so-called formal categories the frequencies of bifaces are abundant in gravelly layer K-30 although the underlying layer K-29 of the same wadi fill produced very few bifaces. It seems that certain activities were carried in the hilly-forested areas from which the more abraded assemblage of K-30 was derived. The almost total disappearance of bifaces in the later assemblages at 'Ubeidiya is noticeable and as yet

unexplained.

Estimated dates for the fossil-bearing strata of the 'Ubeidiya Formation are between ca. 1.6 - 1.2 Ma. Paleomagnetic analysis of the 'Ubeidiya Formation indicated a reversed polarity suggesting that it predates the Brunhes - Matuyama reversal (Opdyke et al., 1983; Braun et al., 1991; Verosub and Tchernov, 1991). Two short, normal paleomagnetic episodes have been found in the Fi member layers II-33 and II 23-24. Layer II-33 has been assigned to the Cobb Mt. (1.215 - 1.190 Ma) and layer II 23-24 has been assigned to the Gilsa (1.575-1.567 Ma) (Sagi, 2005). This correlation fits well with an ESR date of ca. 1.2 Ma for the stratum I 26 which immediately overlays stratum II 33 (Rink et al., 2007).

The dating of these short polarity events was also corroborated by local faunal turnovers (Belmaker, 2009). The 'Ubeidiya fauna (Table 1) can be assigned to a local mammalian fauna biozone older than the assemblages of Bitzat Ruhama, Evron-Quarry and Latamne (Belmaker, 2009). All these sites have been dated to ca. 1.0-1.2 Ma suggesting that the 'Ubeidiya normal polarity events in strata II 23-24 and II-33 should both predate the Jaramillo (0.99-1.07 Ma). Further corroboration by long-range biochronological correlations indicates that the large mammalian assemblage of 'Ubeidiya is similar to the Farneta faunal unit (the sites of Selvella and Pietrafitta, Italy) (Belmaker, 2006; Martínez-Navarro et al., 2009), which has been dated to ca. 1.6-1.2 Ma (Caloi and Palombo, 1997 and references therein). Employing similar approach of long distance comparisons to the lithics of 'Ubeidiya suggests that the assemblages are similar to those from upper Bed II at Olduvai dated to ca. 1.53-1.27 Ma (Bar-Yosef and Goren-Inbar, 1993; Cerling and Hay, 1986).

328

329 **Evron-Quarry**

330 The excavations at Evron-Quarry, located in the coastal plain of the Western
331 Galilee, exposed a sequence of alternating deposits of sandstone (*kurkar*), sometimes up
332 to three meters thick, and red-brown loams (*hamra*), either as isolated lenses or layers up
333 to about one to four m thick (Ronen, 1991b). The layer that contained the Acheulian
334 industry had two distinct horizons of calcareous concretions, occasional artifacts, and
335 sandy clay lenses with pebbles (two to three meter thick) separated from the dark brown-
336 black clay (two m thick) that contained Late Acheulian artifacts and a few animal bones.
337 The artifacts within the archaeological horizon had a vertical distribution of 15-25 cm
338 deep, a phenomenon interpreted as the result of repeated occupations.

339 The archaeological assemblage contained small pebbles of quartz, limestone, and
340 flint, with most of the artifacts made of the latter raw material. No bifaces were found in
341 the excavated areas probably due to their small surface but earlier searches in the quarry
342 dumps recovered twenty handaxes. These are of large size (140-220 mm in length) and
343 demonstrate a relatively crude workmanship that resembles that of the Latamne site
344 (Ronen, 1991b). Large cobbles from which the bifaces were made were recovered
345 together with a group of hard calcite geodes, the heaviest of which was 580g, were
346 brought to the site by the occupants from about 5 km away.

347 Chronologically, the context of Evron-Quarry is assigned to a post-‘Ubeidiya age,
348 and is perhaps contemporaneous with the Latamne site in Syria (ca 1.0-0.8 Ma).
349 Paleomagnetic and ESR studies of the archaeological bearing strata have suggested a date
350 ca. 1.0 Ma (Porat and Ronen, 2002; Ron et al., 2003).

351

352 **Bizat Ruhama**

353 The site of Bizat Ruhama is situated in the southern coastal plain and the
354 archaeological horizon is incorporated in a paludine deposit (Ronen et al., 1998; Zaidner,
355 2003a, b; Zaidner et al., 2003). The retrieved rich lithic industry, generally made of small
356 flint pebbles, falls within the category of “core and flake” assemblage with intensively
357 retouched flakes (small borers, notches and denticulates, etc.) No bifaces were found.
358 Paleomagnetic information and TL dates places the site at 0.99 to 0.85 Ma (Laukhin et
359 al., 2001; Ron and Gvirtzman, 2001).

360

361 **Latamne**

362 The site of Latamne was excavated by D. Clark (1967) and additional fieldwork
363 was conducted by Sanlaville and his associates (1993). The archaeological horizon of
364 Latamne lay in the mid-sequence of the Latamne Formation. It contained the Latamne
365 “occupation floor,” a silt layer only a few centimeters thick (up to 10cm) capped by
366 sandy-silty bedding with traces of rootlets. The sequence was interrupted by erosion and
367 was overlaid by a fluvial sandy deposit, capped by a lacustrine layer (Sanlaville et al.,
368 1993).

369 Geomorphological observations indicate that the archaeological horizon of
370 Latamne resulted from a low energy water flow responsible for the deposition of the
371 artifacts and their pattern of spatial distribution. About one-third of the total recorded
372 artifacts, made from local available flint, limestone and basalt, were slightly abraded or
373 fully abraded. Tool classes can be roughly divided into a collection of bifaces, light duty

scrapers, heavy-duty tools, and a few limestone and basalt spheroids. Several handaxes are trihedral picks similar to those found at 'Ubeidiya.

Water activity caused leaching of the sediments as well as diagenesis that destroyed most of the bones. Most of the well-preserved identifiable bones were collected from the gravels underneath the archaeological site (Hooijer, 1962; Guérin et al., 1993). The entire assemblage reflect a great similarity between the faunas from Latamne and 'Ubeidiya. Originally, the date of the site was estimated as ca. 0.7 Ma based on faunal correlations (Guérin et al., 1993) but the presence of the arvicolid *Lagurodon arankae* and typo-technological affinities of the lithic assemblage suggest a date ca. 1.0 Ma (Tchernov, 1994). However, a single TL date of 560 ka for the Latamne Formation led to the conclusion that the site is 700-500 ka years old (Sanlaville, 1988; Sanlaville et al., 1993) which seem untenable.

Dursunlu

The quarry of Dursunlu is 75 km northwest of Konya, western Turkey. It comprises Early Pleistocene lucustrine limestones, marls and clay with lignites. Paleomagnetic analysis has suggested a reverse polarity punctuated by three episodes of normal polarity, consistent with an Early Pleistocene age. A more detailed analysis of the drilled cores suggested that the artifacts were derived from two layers between the Jaramillo and MBB, i.e., 0.99 and 0.78 Ma (Güleç et al., 2009). Most of the small lithic assemblage was made of quartz with rare pieces of flint and an igneous rock. The assemblage falls under the category of core and flake industry and provides a unique point in the presence of early hominins in Anatolia.

397

398 **Gesher Benot Ya'aqov**

399 The site of Gesher Benot Ya'aqov (GBY) lies in the Jordan Valley, at the eastern
400 edge of a vast, basalt-covered area. The excavations in the 1930s by M. Stekelis, and
401 recently by Goren-Inbar and her associates (Goren-Inbar et al., 1991, 1992a, b, 2000,
402 2004; Goren-Inbar and Saragusti, 1996; Alpers-Afil et al., 2007) provided an African-
403 type assemblage of cleavers and bifaces that is unlike any of the other known 200
404 Acheulian occurrences in southwestern Asia including both surface and excavated
405 occurrences (e.g., Hours, 1975, 1981; Bar-Yosef, 1987; Goren-Inbar, 1995; Liubin, 2002;
406 Taskiran, 1998). The nature of the deposits and the malacological assemblages,
407 dominated by *Viviparus apameae*, indicate that the archaeological assemblages
408 accumulated on the shores of an expanding lake that flooded the gorge (Horowitz, 1979;
409 Goren-Inbar et al., 1991, 1992a, b; Goren-Inbar and Saragusti, 1996; Feibel, 2004).

410 The complex stratigraphic sequence, first partially exposed by M. Stekelis (1960)
411 and more recently by systematic excavations conducted by N. Goren-Inbar (Goren-Inbar
412 et al., 2004; Feibel, 2004) contains early layers with Acheulian industry dominated by the
413 production of cleavers and bifaces from basalt, as well as flint and limestone objects.
414 Some of the cleavers were fabricated by the Kumbewa technique. Although lava flows
415 cover the area, on both sides of the parts of Jordan River and in particular the large area
416 of the Hauran-Golan no other Acheulian sites with basalt industry were located. On the
417 contrary, in most cases flint nodules derived from isolated limestone and chalky outcrops,
418 often of Eocene age, served as raw material for fabricating handaxes (e.g., Goren, 1979;
419 Goren-Inbar, 1985; Ohel, 1991).

420 The archaeological horizons of GBY are embedded in a depositional sequence
421 that accumulated above a lava flow with normal polarity. The lava flow, designated as the
422 Yorda Basalt, was first K/Ar dated to 0.68 ± 0.12 Ma (Horowitz, 1979) and later to
423 0.9 ± 0.15 Ma (Goren-Inbar et al., 1992a). However, the paleomagnetic sequence within
424 the excavated deposits demonstrated that the Matuyama Brunhes Boundary (MBB) was
425 located within the series of deposits and the accumulation of the entire sequence took
426 place during ca. 100,000 years (Feibel, 2004).

427

428 **Holon**

429 The site of Holon is embedded in marshy deposits overlying an abraded kurkar
430 ridge dated by Horowitz (1979) to around 500-400 ka. The site contained more than one
431 level (Chazan and Kolska-Horwitz, 2007). ESR and luminescence dating (Porat et al.,
432 2002) suggested for the archaeological horizon an average date of ca 205 ka based on
433 ESR and TL dates and suggested that is comparable to the ESR readings from teeth from
434 Tabun layer E of Garrod's excavations (the Acheulo-Yabrudian industry). However, TL
435 dates of burnt flint pieces from Tabun E indicate an age older of 270 ka, considered as the
436 onset of the Mousterian (Mercier et al., 1995) dated also in other sites around 220-250
437 ka. Unfortunately the dates for Holon were retrieved from another geological exposure as
438 the area of the original site lies below a major factory.

439 The bifaces of Holon are mostly of pointed and rounded aspects; the flake
440 industry contains side scrapers, denticulates, and notches, along with cores and debitage
441 products This lithic assemblage has been attributed to the Late Acheulean (Chazan and
442 Kolska-Horwitz, 2007) and if the dates are accepted it would mean that either the Holon

Acheulian was contemporary with the early Mousterian, which is untenable, or that there is a systematic difference between the luminescence dates of quartz grains that were retrieved from another location although near the location of the original excavation.

Qesem

Qesem Cave is located 12 km east of Tel Aviv, Israel. Excavation in the cave revealed several meters of deposits with abundant lithic and faunal assemblages and plenty of evidence for systematic use of fire (Karkans et al., 2007). Uranium series dates have suggested an age bracket of 0.4 - 0.2 Ma (Barkai et al., 2003). The rich lithic assemblages, dominated by the production of blades, were assigned to the Amudian culture (Gopher et al., 2005; Barkai et al., 2006; Lemorini et al., 2006).

Revadim

Revadim Quarry is located on the southern coastal plain of Israel, north of Kibbutz Revadim. Paleomagnetic dates suggest a normal polarity indicating an age younger than 0.78 Ma. OSL dates suggest a minimum age bracket of 0.3-0.2 Ma. (Gvirtzman et al., 1999; Marder et al., 1999). The lithic assemblage at the site suggests a high frequency of flake tools and the presence of handaxes. The typo-technological characteristics of the bifacial tools have suggested that it may be attributed to the Late Acheulean culture (Marder et al., 1999, 2006).

Oumm Zinat

The site of Oumm Zinat is situated 500 m east of Kibbutz Evron and was first excavated by Prof. Stekelis in 1950. Subsequent excavations by Gilead and Ronen in 1977 revealed a small to medium size assemblage dominated by hand axes contemporaneous with Late Acheulean such as Oum Qatafa D1 and Tabun E/F (Kolska-Horwitz and Tchernov, 1989)

Tabun E

Tabun Cave is located on the Southwest flanks of Mount Carmel, Israel on the banks of Wadi el Mughara (Nahal Me'arot). The excavation of the cave began in 1930's under Bate and Garrod and continued with Jelinek (1970-1972) and recently A. Ronen since the 1990's. Layer E consists of slightly abraded flint over an area of 1.0X1.3 m. The lithic assemblage was characterized as Late Acheulean (Garrod and Bate, 1937).

The TL dates for Tabun E to D suggested that they were deposited during MOIS 9 through 8 (Mercier et al., 1995), more recent dates by Mercier and colleagues (2000) suggest that Tabun F and E were deposited during MOIS 9 between 350 ± 30 and 330 ± 30 ka. However, ESR dates for Tabun E by Grün et al., (1991) and Grün and Stinger (2000) suggested a date of only 200 ka.

Oumm Qatafa

Oumm Qatafa is situated on the left bank of Wadi Khareteitun in the Judean Desert, ca. 15 km west of the Dead Sea. The site was excavated by R. Neuville between 1928-1949 (Neuville, 1951) where he identified a sequence which spans the chacholithic

through the Lower Paleolithic. The Upper Acheulian level in D2 is characterized by broad, ovate discoidal and cordiform bifaces. Recent ESR dates obtained from level D2 indicate an upper date of 213 ± 26 ka and thus an end for the late Lower Paleolithic (Porat et al., 2002).

Early and Late Acheulian in the Levant

Surveys along the Levantine coast located a few occurrences that appear to be of Early Pleistocene age although dating, in most cases, is rather tenuous due to lack of datable substances. Shorelines were dated on the basis of their elevation above sea level, while the known relative ages of foraminifera and marine shell assemblages were also incorporated into these figures along the western mountainous range along the Levantine coast, a few artifacts were found on terraces as high as 120 m above sea level. We therefore choose to summarize those sites where systematic excavations or surveys were conducted and their dates depend of the geological stratigraphy and its suggested correlation with sea level chronology. Here we add some brief information about the main occurrences, and for a full survey see the available summaries (e.g., Hours 1975, 1981; Muhesen, 1985, 1993; Sanlaville et al., 1993; Bar-Yosef, 1998a).

One of the distinctive sites is Kefar Menachem, situated in the interior part of the coastal plain and was excavated twice (Gilead and Israel, 1975; Barzilai, 2006). The lithic industry is embedded in red loam of an Early Pleistocene age (Horovitz, 1979). The lithic assemblages of both excavated areas are comprised of numerous core and flake products, a few flake-tools (classified as end-scrapers, side scrapers, burins, notches, and denticulates). The use of direct hard hammer percussion is dominant. To date, the rare

bifaces have been found only on the surface and their attribution to the excavated collections is doubtful. These bifaces are described as irregular ovates, picks, long lanceolates, and backed bifaces (Gilead and Israel, 1975) who tentatively related this surface assemblage to the Early Acheulian.

Most of what is known from the areas of Lebanon and Syria was obtained through the study of the terraces of Nahr el Kebir, the Orontes, and the Middle Euphrates. The majority of the collections were classified as Early and “Middle Acheulian” first on the basis of stratigraphic grounds and later upon consideration of their typological characteristics. Rare finds were retrieved and in several localities in the Beqa’a Valley.

Lithic studies of the so-called “Middle Acheulian” assemblages identified two geographic facies. The sites along the coast, such as Berzine and Ouadi Aabet, contain essentially amygdaloid and oval bifaces, while the inland sites (Joub Jannine II and Latamne) have more lanceolates and trihedral picks, and along the Nizip river (Minzoni-Déroche and Sanlaville, 1988), a tributary of the Euphrates in Turkey, recognized a similar distribution of the “Middle Acheulian” artifacts.

The general technological tendency among the Acheulian sites is toward a greater use of soft hammer percussion and the sporadic appearance of the Levallois technique (Copeland and Hours, 1981). Typologically, the almost total disappearance of core-choppers is noticeable. The cordiform and amygdaloid bifaces outnumber the ovates. The length of the handaxes decreases in general, a tendency that was already noted by D. Gilead (1970) for the southern Levantine samples.

The sites of Hummal and Nadaouiyeh I were discovered in El-Kowm basin (Hours et al., 1983; Le Tensorer et al., 1993), and exposed several occupational horizons.

Nadaouiye h I comprised of an accumulation of alternating clayey layers and by sandy layers near an artesian spring. The stratigraphy is a complex one, marked by slumping and erosion. The systematic excavations demonstrated the presence of *in situ* Acheulian assemblages. The bifaces, generally amygdaloid, were accompanied by a rich flake industry in every one of the six tested layers. The presence of the Levallois technique was noted in low frequencies. In addition, the both Hummal where a much longer depositional sequence was revealed and Nadaouiye h contain Yabrudian or Acheiulo-Yabrudian capped by Hummalian (an early Mousterian industry) contexts (LeTensorer 2004).

Subdividing the Late (or Upper Acheulian) into phases or “facies” was and still is not an easy task. On the basis of technological and morphological considerations Gilead (1970) subdivided the Late Acheulian into several groups as follows:

A. The Ma’ayan Barukh group (MB), mostly from open-air sites, is characterized by the dominance of the cordiform aspect (including amygdaloids, cordiforms and subtriangulars) with fewer ovate and a few pointed bifaces and rare cleavers. The assemblage from the excavations of Umm Qatafa D2 is included in this group.

B. The Evron-Kissufim group (EK) is, on the basis of stratigraphic evidence, later than the MB group. It contains a richer flake tool component, up to 30-60%, with clear evidence for the manipulation of the Levallois technique. The bifaces show a decrease in rounded aspects (ovates and discoids) and a slight increase in the pointed forms.

C. The Sahel el-Khoussin-Yiron group (SY) are those assemblages mostly surface collected in the hilly areas and flanks. The bifaces are somewhat cruder than those of the other groups with an occasional dominance of the rounded aspect over the cordiform

aspect (Yiron, Beith Uziel, Baqaa-Rafaim etc.). As in the EK group, the Levallois technique was practiced in some sites. It is worth noting that despite the hilly distribution, these assemblages are not present in the three caves where Late Acheulian layers were uncovered (Tabun F, Abu Sif, Umm Qatafa D).

The frequency of refinement index (thickness/breadth x 100) demonstrates the differences among the sites. The same is probably true when the mean length among Late Acheulian sites is considered. Wherever large cobbles were available, there was a tendency towards larger bifaces. However, a general tendency for decrease in biface length could indicate increasing efficiency of resharpening (perhaps longer curation?) during the Late Acheulian.

The flake industry of most of the Late Acheulian occurrences is not very well known. In some places, the number of flakes cannot account for their manufacture. For instance, the thousands of bifaces found in Ma'ayan Barukh may have been produced in an area further north near the Litani River. The flakes collected from the same surface clusters could indicate some resharpening (although small flakes and chips are not easy to retrieve in the deep red soil of these hills). It seems that the concentration of bifaces near the Hula Lake shores on the interfluves of freshwater creeks may represent repeated butchering activities in a pristine environment.

A unique Late Acheulian site, embedded between a lava flow dated to 233 ± 3 ka and an older lava flow dated to 800 ka, was excavated on the edge of the crater lake known as Berekhat Ram on the Golan plateau (Goren-Inbar, 1985). The rich assemblage contains several thousands of artifacts, mostly in mint condition, with about four hundred of retouched pieces including eight small bifaces. The makers of the industry employed

the Levallois centripetal (radial) technique. A special find is a human figurine (Goren-Inbar, 1986) that recently received much attention in the debate concerning the capacities of Archaic *Homo sapiens* or late *Homo erectus* (e.g., Marshack, 1997; D'Errico, 2000). The actual date of the site is unknown but given the current TL dates of the Acheulo-Yabrudian, it should be placed during the time span of 350-450 ka.

In general, Late Acheulian sites can be found across the southwestern Asia in every environment including the coastal plain, hilly areas, inter-montane valleys, and oases and in desert landscapes. The best example to date from an oasis situation is the series of Late Acheulian assemblages which is characterized by high frequencies of bifacial cleavers, uncovered in the Azraq basin (Copeland and Hours, 1989; Rollefson, 1997). Among these, the sounding at Lion Spring provided stratified lithic assemblages that are characterized by ovate, amygdaloid, and cordiform bifaces, with a rich flake industry. In the absence of precise dating and on the basis of comparisons with the occupations in other location it seems that some Late Acheulian occupations should be correlated with periods of wetter conditions during the time span of ca. 600-400/350 ka.

The Acheulo-Yabrudian

The Acheulo-Yabrudian, was renamed as the Mugharan Tradition by Jelinek (1981, 1982a, b), and has a definite geographic distribution from the Taurus foothills to the central Levant (Bar-Yosef, 1998b). It is generally dated to 400/350- 250/220 Ka, and is found stratigraphically always under the Early Mousterian (e.g., Tabun, Hayonim, Misliyah caves; Garrod and Bate, 1937; Weinstein-Evron et al., 2003). We stress that in spite of intensive surveys, none of the typical artifacts by which this entity is defined

were found in the Negev and Sinai or in the desert region of southern Jordan. Three facies that sometimes considered as independent industries were identified on the basis of quantitative studies, as follows (Jelinek, 1982a, b; Copeland and Hours, 1983):

1. The Yabrudian facies contains numerous side-scrapers, often made on thick flakes including canted ones, thus resulting in relatively high frequencies of Quina and demi-Quina retouch with a few Late Paleolithic tools and rare blades. Although typologically, Levallois-type products have sometimes been mentioned, the reconstruction of operational sequences has not yielded a well-identified Levallois method.

2. The Acheulian facies is considered by Jelinek (1982a, b) to consist of up to 15 percent bifaces, with numerous scrapers fashioned in the same manner (Quina type scalar retouch) as the Yabrudian ones.

3. The Amudian facies is characterized by end scrapers, burins, backed knives, and rare bifaces and was therefore originally called 'Pre-Aurignacian' in the sense of being blade dominated. Not surprisingly, until the 1940s the term Aurignacian was used to refer to all early Upper Paleolithic industries in Western Eurasia. The Amudian facies, following the Tabun excavations, seems to be closer typologically to the Acheulian than to the Yabrudian and contains evidence for limited practice of the Levallois technique (Jelinek, 1982a, b). The "Pre-Aurignacian" in Yabrud I and Abri Zumoffen are richer in "Upper Paleolithic" elements. The excavations at Qesem cave produced rich assemblages of blades and plenty of evidence for the use of fire (Gopher et al., 2005; Barkai et al., 2006; Karkans et al., 2007) and a faunal assemblage dominated by Fallow deer (Stiner et al., 2004).

625

626 **The Lower Paleolithic of Iraq, Iran and the Caucasian region**

627 The vast geographic area summarized here is generally poorly known. The scant
628 evidence from Turkey on one end of the region and India on the other, including a few
629 recorded find spots from Iran (Smith, 1986) and from the Arabian Peninsula (Zarins et
630 al., 1979, 1980, 1982; Whalen et al., 1983, 1984; Abdul Nayeem, 1990), indicate that
631 bifaces can be found everywhere. The distribution toward the northern edges of the Near
632 East has implications for the reexamination of the “Movius line”.

633 In Iraq, little is known beyond the site of Barda Balka, located in the Chemchemal
634 valley in Kurdistan, which was collected and excavated by Howe (Braidwood and Howe,
635 1960). This predominantly flake assemblage may be of Middle Paleolithic age. Iranian
636 finds are also few and are far apart. In Khorasan, on the edge of a dried-up lake, quartzite
637 and andesite core-choppers were collected (Ariai and Thibault, 1975/77). In the absence
638 of dates, the investigators related the assemblage to the Late Pliocene on typological
639 grounds. Isolated bifaces have been collected in various places in Iran (Smith, 1986). The
640 Ladizian industry in Baluchistan (Hume, 1976) to the east should be mentioned briefly. It
641 is defined on the basis of scatters of lithics on old river terraces and it is a core-chopper
642 industry with retouched pieces but no bifaces. Hume (1976) has proposed a late Middle
643 Pleistocene age for the Ladizian.

644 Early and Late Acheulian sites in Georgia were either surface collected or
645 excavated (Liubin and Bosinski, 1995; Liubin, 2002). Among the surface sites, some of
646 the interesting collections were done in Cikiani, near Paravani in southern Georgia, where
647 cleavers and handaxes were found (Kikodze, 1986). The bifaces were made of andesite

but most of the cores were made of obsidian. Persati, another surface site, which is yet unpublished, is located on top of a volcanic plateau that is apparently the continuation of the Dzavacheti range in southern Georgia, about 2100m above sea level (Kikodze, personal communication). The artifacts were found at the edge of lacustrine sediments, dated to the Neogene. One of the find spots seems to be eroding from these lake deposits, but radiometric dates are not available.

Acheulian industries were uncovered in four excavated caves: Azych, Kudaro I and III, and Tsona. The lithic industry in Azych, located 800 m above sea level, is subdivided on the basis of the observed stratigraphy into several phases of the Acheulian. The earliest levels produced a few core-choppers but they are without clear attribution to a prehistoric entity (Liubin and Boskinski, 1995; Liubin 2002). The richest assemblages were uncovered in layers VI and V, including mainly Late Acheulian bifaces with distinct use of the Levallois technique. In layer V, a fragment of a hominid mandible was found.

Kudaro I is situated 1600 m above sea level. Layer 5 contained Acheulian artifacts made from local raw material. The entire assemblage is characterized by a high frequency of retouched pieces including numerous side scrapers. Core-choppers and bifaces, mostly of elongated shapes, assign this assemblage to the Late Acheulian. The presence of flake cleavers was noted by the excavator. Three human teeth were also found in this context. Dates suggest a range of 250-300 ka. TL readings indicate a slightly earlier time, 360 ± 90 Ka and 350 ± 70 ka. However, a reversed paleomagnetic situation identified in level 6 immediately below the Acheulian, with the fauna of several Galerian elements, hints that the Acheulian in layer 5 is perhaps of an older age (Liubin and Bosinski, 1995).

Kudaro III produced a more restricted Acheulian collection in layers 6-8 with bifaces and flake tools. A TL date of level 8a was 560 ± 112 ka, while layer 5 produced dates of 252 ± 51 ka and 245 ± 49 ka (Liubin, 2002).

Tsona Cave, at an altitude of 2150 m above sea level, and just about 5-6 km south of Kudaro, is a very large cave. The Acheulian industries were derived from layers 6-7a. Preliminary reports indicate that the lower assemblages (layer 7) that were originally considered as Early Acheulian produced only a small sample. The assemblage of layer 6 assigned to the Late Acheulian contained about 100 artifacts, including 29 bifaces made from local raw material, mostly retrieved in the form of pebbles. The chronological position of this Acheulian is not well-known, however it seems that this cave, like others at high altitudes, could have been occupied only during interglacial times and in most cases were possibly only seasonal hunting camps. At least two other cave sites on the northern flanks of the Caucasus are known to contain Late Acheulian remains.

In sum, the distribution of the Acheulian industries in Western Asia is essentially limited mainly to the Levant with decreasing abundance in the Caucasian region and eastern Anatolia. In spite of the paucity of information from Iran, as mentioned above, there is no evidence for continued distribution of the Acheulian into India. Thus, the “Movius Line,” which distinguishes between the Acheulian and the non-biface industries, separates Anatolia from Eastern Europe and the Iranian plateau and Baluchistan from the Levant.

Human remains

While the lithic stratigraphy in the region is rather complete, the hominin remains

are woefully underrepresented. The Lower Paleolithic sequence of Western Asia, apart from Dmanisi, is very poor in human fossils. Dmanisi, as mentioned above, contained several skulls and postcranial bones, demonstrating a large range of morphological variability (Gabunia and Vekua, 1995; Braüer and Schultz, 1996; Gabunia et al., 2000a; Gabunia et al., 2002; Lordkipanidze et al., 2007; Rightmire et al., 2006).

The available human remains from this long period are scanty, and a few are surface finds. The extensive excavations at ‘Ubeidiya (1959-1999) have revealed numerous lithics which attest to hominid presence in the Southern Levant during the Early Pleistocene. Several cranial fragments (UB 1703, 1704, 1705 and 1706), an incisor (UB 1700) and a molar (UB 1701) have been assigned to *Homo* indet. (Tobias, 1966a, 1966b) and as *Homo cf. erectus* (Tchernov and Volokita, 1986). Belmaker et al., (2002) described a new right lateral hominid incisor (UB 335) from stratum I-26a and which was assigns to *Homo cf. eragster*. However, the antiquity of the previous finds has been contested, and UB 335 is the only one that can be securely assigned to the Early Pleistocene deposits.

Two broken femora from GBY (Geraads and Tchernov, 1983), attributed to *Homo erectus*, were identified in the collections of animal bones made at the site when the deepening of the Jordan River channel took place.

A broken femur was uncovered in Tabun cave layer E (McCown and Keith, 1939) within the Acheulo-Yabrudian assemblage. It thus occupies the same stratigraphic and chronological position as the fragmentary skull from Zuttiyeh (Gisis and Bar-Yosef, 1974). The latter is considered as an example of an Archaic *Homo sapiens* (Vandermeersch, 1995) and could have been one of the potential ancestors of the later

Qafzeh-Skhul group. Recently, this fragmentary skull has been compared to the Zhoukoudian human remains and interpreted as belonging to a generalized Middle Pleistocene Asian population (Sohn and Wolpoff, 1993).

Only after MOIS 5 do hominin remains (both of Anatomically modern humans and Neanderthals) become more common in the region and allow to fully discuss aspect of biological adaptability vis á vis both environment and culture.

Early-Middle Pleistocene Environmental changes

The unique geological-climatological position of the Levant in conjunction with a highly fragmented habitat contributes to creating an extremely varied and complex biotic picture. The great diversity of the Southern Asian biota, both in number of species and in biogeographic origin (Palearctic, Palearctic and Saharo-Arabian), is primarily the product of the Afro-Eurasian biotic interchanges over the past 25 million years (Miocene through Quaternary).

Since the beginning of the Miocene Southwestern Asia has been situated at the crossroad between African and the Eurasian continent thus forming a wide land bridge between the two regions. During different time periods, tectonic, climatic and sea level changes altered the “permeability” of this land bridge. At times the region would allow for animals to disperse in opposite directions and in other times the land bridge served as a barrier. The Messinian event at the end of the Miocene represents a period of long distance reciprocal biotic exchange due to the dramatic lowering of the Mediterranean Sea level. However, since then the developing Red Sea finally limited this geographic

corridor to the Levant and the southern Arabian Peninsula. Indeed, these two passages functioned as a selective filter for most organisms, through which only specific species could spread or pass (Tchernov, 1988; Tchernov and Belmaker, 2004).

The onset of the Pliocene is marked by an abrupt transgression of the Mediterranean and reestablishment of the barrier between Africa and Eurasia. Thus, Africa became considerably isolated from the rest of the world by the Saharo-Arabian arid belt. North Africa was even further isolated from both sub-Saharan Africa and the eastern Mediterranean region (Thomas, 1985). During this period, the main biotic influences in the region were Asiatic, primarily from the Irano-Turanic region. These are recorded mainly in Anatolian sites, which include many forest dwellers such as the Cervids. The Taurus-Zagros Mountains and the Saharo-Arabian developing desert belt eventually became a biogeographic barrier, particularly for laurophyllous (evergreen broad-leaved) plant species (Jacobs et al., 1999), forest dwellers (Cervidae, Castoridae, Gliridae and Ursidae) and aquatic animals. Hence, the transgression of the Mediterranean increased the isolation of the Southwestern Asia

Towards the end of the Pliocene and in the Early Pleistocene, the regression of the Mediterranean terminated the quasi-isolation of the region and the increase glacial-interglacial cycles had their impact on the region under discussion.

Vegetation and pollen

The bimodal Mediterranean climate has been shown to appear in Southwest Asia during the late Pliocene ca. 3.2 Ma as part of a global cooling trend and was fully established by 2.8 Ma (Suc, 1984). Unfortunately, long pollen cores that would allow for

763 a continuous paleobotanical reconstruction throughout the Early and Middle Pleistocene
764 are rare in Western Asia in comparison to marine cores in central from the Western
765 Mediterranean Sea and the Aegean. Nonetheless, these provide a scale to which we can
766 compare results from smaller cores obtained in the region.

767 A succession scheme has been developed for the Mediterranean vegetation for
768 glacial- interglacial cycles such that glacial steppe vegetation in replaced by
769 sclerophyllous woodland (dominated by *Quercus*, *Pinus*, *Betula* and *Juniperus*) and then
770 by deciduous forests (deciduous *Quercus*, *Carpinus*, *Ostrya* and later on *Abies* and
771 *Fagus*), followed by a regressive phase of open woodland. Specific regional pattern
772 reflects local climatic conditions but pollen analysis in the Mediterranean basin has
773 shown that the vegetation succession follows the processional insolation cycles
774 (Tzedakis, 2007). Thus during the Early Pleistocene, while the glacial- interglacial cycles
775 were dominated by the obliquity 41 ka cycles, the succession of floral communities
776 followed a cycle roughly every 23 ka.

777 A long pollen core in Teneghi Philippon, Greece, that spans the last 1.35 million
778 years, allows us to test general trends in vegetation in relation to orbital forcing. Flora
779 reconstruction for the glacial periods were based on the minima of the AP percentage and
780 are reflected during MOIS 22, 16, 12, 10, 8 and 6 with 16 (0.65 Ma) being the most
781 extensive during the past 1.35 Ma. In early periods such as MOIS 38, 40 and 42,
782 *Artemisia* levels were over 90% indicating extreme aridity, however these conditions
783 were not sustained over a period longer than 10ka (Tzedakis et al., 2006).

784 The majority of pollen data is derived from late Pleistocene and Holocene
785 sediments and post date the period discussed in this study. A playnostraigraphy of the

Jordan Valley (Horowitz, 2001) provides a generalized Quaternary sequence for the Levant, but unfortunately is not well dated. The sequence records alternations between wet and dry Mediterranean flora. The interglacial flora was poor in arboreal pollen in comparison to modern day conditions. A north – south gradient indicated 3- 5% arboreal pollen in the north associated with steppe elements while southern cores indicate no arboreal pollen but proliferation of desert plants. In comparison, the pluvial periods or the glacial flora was dominated by arboreal pollen, and of interest is the increase of pollen the winter deciduous oak *Quercus ithaburensis* (Horowitz, 1988).

Pollen from archaeological sites, in spite of the time gaps, provides another angle to landscape reconstructions, and the conditions that faced the early hominins. A study in Dmanisi indicated a forested area with *Abies*, *Pinus*, *Fagus*, *Alnus*, *Castanea*, *Tilia*, *Betula*, *Caprinus* and rarely *Ulmus* and *Salix*. Bushes and shrubs are represented by rhododendron, *Corylus* and myrtle and the herbaceous vegetation is dominated by Cyperaceae, Germinae and Polygonaceae (Dzaparidze et al., 1992) consistent with the humid thermophilous broad leaf forest zone which is similar in the region today in the western Caucasus (Zazanashvili et al., 2000).

Only few botanical remains have been retrieved from ‘Ubeidiya, all from the Li cycle. Macrofloral remains of fossilized leaves were discovered in the limnic laminated layer III- 19. These have been identified as *Pistacia lentiscus*, *Rhus tripartita* and *Myriophyllum* sp. (Lorch, 1966). Pollen spectrum was extracted from layer III-12, analyzed by A. Horowitz (Bar-Yosef and Tchernov, 1972) indicating 82% arboreal species of which the overwhelming majorities are *Quercus* sp. followed by *Juniperus* sp. and *Olea* sp. Non-arboreal families include Gramineae, Cruciferae and Compositae.

809 Cyperaceae comprise 8.4% of the pollen and attest to the water habitat present at the site
810 suggesting a period more humid than today.

811 There is a small pollen spectrum from Bitzat Ruhama ($n=114$) reflecting
812 relatively high frequencies of arboreal pollen including *Quercus*, *Pinus*, *Olea* and *Cedrus*.
813 (Zaidner, 2003b). Among the non-arboreal pollen we encounter *Chenopodiaceae*, *Palmae*,
814 *Poaceae* and *Liliaceae*. The presence of cedar is indicative of a colder environment in the
815 region than the present (Zaidner, 2003b).

816 A pollen sample from the Latamne formation indicates that mountain slopes near
817 the site were forested by board leaf trees such as *Quercus*, *Caprinius*, *Tilia*, *Juglans*,
818 *Ulmuss*, *Corylus* and *Betula* and coniferous species such as *Pinus* and *Cupressus*
819 (Dodonov et al., 1993).

820 The wood remains of GBY suggest the presence of Mediterranean wood and plant
821 species and that the climate in the Hula valley at the time of deposition resembled the
822 seasonal Mediterranean pattern seen today. Among the dominant species are oak
823 (*Quercus* sp.), wild pistachio (*Pistacia atlantica*), wild olive (*Olea europaea*), plum
824 (*Prunus* sp.), and jujube (*Ziziphus spina-christi*) (Goren-Inbar et al., 2004; Werker, 2006).
825 Immersed or floating freshwater plants are common at the site and include species such as
826 *Euryaleferox*, *Najas foveolata*, *Nymphoides* cf. *peltata*, *Potamogeton*
827 *coloratus/polygonifolius*, the extinct *Stratiotes intermedius*, *Trapa natans* and apparently
828 *Sagittaria sagittifolia*. Bank forest taxa include wild grape (*Vitis sylvestris*) and ash
829 (*Fraxinus syriaca*).

830 Pollen analysis from this site (van Zeist and Bottema, 2009) supports the general
831 landscape reconstruction derived from wood identification specifically the presence of

832 *Quercus ithaburensis* – *Pistacia atlantica* woodland land or park-forest. Above 400-
833 500m to the west of the valley, this Tabor Oak woodland species was replaced by
834 *Pistacia atlantica*- *Amgdalus korschinskii* woodland. The overall pattern indicates that
835 the upper Jordan Valley did not undergo any significant vegetation changes as a results of
836 the glacial- interglacial periods as we see in higher latitudes and the overall vegetation as
837 observed in the wood remains as in the pollen reflects comparable environs to those of
838 today. However, oxygen and carbon stable isotope evidence derived from gastropods and
839 ostracods at GBY as well as evidence from mollusk ecology indicate a climatic shift
840 (Rosenfeld et al., 2004; Ashkenazi et al., 2009; Spiro et al., 2009) inline with global
841 climatic forcing throughout the sequence across the MBB, the paleoecological
842 reconstructions based on vegetations (pollen, wood) indicated that climatic forcing may
843 have had a stronger effect on the local hydrological regime and aquatic fauna than on the
844 terrestrial ecosystem and most notably fauna (see Kingston, 2007, for further discussion
845 of this phenomenon).

846 847 *Faunal turnovers and dispersal from Africa*

848 The paucity of sites from the Early and Middle Pleistocene does not allow to
849 track both inter and intra variability within each of the sub-regions during this long
850 period. There are far less sites that preserve well-dated faunal assemblages than lithic
851 localities that can be used for paleontological and paleoecological analysis. The
852 observations made on the basis of fossiliferous sites are generally correlated with the
853 European and South African sequences (Belmaker, 2009). While the actual species
854 composition within each region is different the tempo and mode of change are similar

across the region and point to a secular trend of cooling through time. Data for this analysis is derived from archaeological sites (see previous section for detailed description of the sites) as well as a several paleontological site that yielded only faunal material.

The Late Pliocene is well recorded in the Caucasus such in Kvabebi, (Georgia) which yielded a large assemblages indicative of a marshy and riverine habitat (Vekua, 1995) surrounded by open woodland environment (Agustí et al, 2009b). Similarly, the Bethlehem fauna with the small assemblage of 11 mammalian species can be identified as representing a humid and temperate woodland environment (Gardner and Bate, 1937; Hooijer, 1958). The presence of *Giraffa* sp. uncovered in Bethlehem suggests an African dispersal during the Late Pliocene although the poor preservation of the specimen precludes identification to species (Robinson and Belmaker, in press). The presence of the three-toed horse *Hipparion* (Hooijer, 1958) would indicate a date ca. 2.5 Ma. During this time period, the Jordan rift had not formed in its current state (Horowitz, 2001), which may have hindered a large dispersal from Africa. Furthermore, as previously mentioned, the absence of sites prior to Bethlehem does not allow us to pinpoint the precise date of this dispersal which may have occurred any time prior to that.

The transition from the Late Pliocene to Early Pleistocene, similarly to the Olduvai dry event, indicates a shift from a more humid and wet environment to a more open, partially dry and somewhat cooler conditions. The change is attested when the faunal assemblages of Bethlehem and Kabevebi are compared to Dmanisi (1.8 – 1.77 Ma) (Gabunia et al., 2000; Agustí et al, 2009b), Kotsakhuri (1.63 – 1.83 Ma) and Tsalka (Vekua and Lordkipandize, 2008). While the fauna from these sites is characteristic of Mediterranean temperate woodland indicated by a high proportion of cervids, an

ecological shift to a more open habitat is marked by the first appearance of several taxa such as the Mammoth (*Mammuthus meridionalis*) and the large Stenoid horse (*Equus stenonis*).

A local faunal turnover can be observed between the Early Pleistocene fauna dated to ca. 1.8 – 1.7 Ma as represented by Dmanisi, Kostakhuri and Tsalka, and the younger site of ‘Ubeidiya (1.6 – 1.2 Ma). While all sites have many taxa in common, the older sites exhibit a more archaic suite of species including *Canis etruscus*, *Cervus perrieri* and *Eucladoceros* sp. replaced by the younger *Canis mosbachensis* and late Villafranchian cervids *Praemegaceros obscurus- verticornis* at ‘Ubeidiya. ‘Ubeidiya possibly represents a long sequence spanning ca. 400 Ka and its landscape was essentially Mediterranean woodland similar to that of Dmanisi as shown by the great abundance of cervids and other Eurasian taxa (Belmaker, 2009). However, several new taxa that appear in ‘Ubeidiya are not present in the previous Late Pliocene deposit of Bethlehem (ca. 2.5 Ma) and the Early Pleistocene sites in the Caucasus. These include the unique suite of African taxa such as *Pelorovis oldwayensis*, *Kolpochoerus olduvaiensis* and Cercopithecinae cf. *Theropithecus* sp. (Belmaker, 2010), which coincides with the Aullan dispersal event dated to 1.8 Ma (Arribas and Palmqvist, 1999). This indicates that a dispersal route from East Africa to Eurasia became available during this time or immediately prior to that date.

The comparison between ‘Ubeidiya and Dmanisi is particularly informative. With the exception of the genus *Homo*, the Early Pleistocene sites of the Caucasus are devoid of African taxa (Tappen, 2009 but see Martínez-Navarro and Palmqvist, 1995), and have a high proportion of endemic species such as *Bison georgicus* and *Struthio dmanisiensis*.

This suggests that during the Late Pliocene the Caucasus was isolated and this situation allowed for the evolution of endemic species. However, ‘Ubeidiya and the Georgian sites share several taxa and specifically the jird, *Parameriones obiediensis*. This species was identified as an endemic to ‘Ubeidiya and subsequently found in Dmanisi indicating that the biogeographic route connecting the two regions was open. In addition the path from East Africa into the Levant was also open and led to the influx of African taxa into the latter.

The Epivillafranchian fauna is represented by the sites of Bizat Ruhama, Evron Quarry and Daquara, Latamne in the Levant, Dursunlu in Anatolia and Akhalkalaki in Georgia. The archaic nature of this fauna is maintained by the presence *Hipopotamus behemoth* in Latamne. However, a novel dispersal from Africa is attested by the arrival of the suid *Kolpochoerus evronensis* in Evron Quarry and by *Giraffa camelopardalis* at Latamne (Robinson and Belmaker, in press). Furthermore, the appearance of modern taxa such as *Dama mesopotamica* and *Cervus elaphus* indicates a new and more modern faunal composition.

The increased shift towards a more open, arid habitat, compared to earlier sites dominated by cervids and woodland taxa, is evidenced by the increasing dominance of equids and the steppe mammoth (*Mammuthus trogontherii*). The small faunal assemblage of Bizat Ruhama (ca. 1.0 Ma) in the southern coastal plain has only bovid, equid and hippo remains (Ronen et al., 1998). Evron - Quarry (ca. 1.0 Ma) although situated at the western margins of the Galilee where rainfall is higher than in the south, the cervid sample comprises only of four among a total of 36 (11%) identified specimens, while bovids are represented only by 11 specimens (30%) (Tchernov et al., 1994).

924 Akhalkalaki (0.9-0.8 Ma) in central Georgia is dominated by megacerids,
925 *Stephanorhinus etruscus*, *Mammuthus trogontherii*, *Equus suessenbornensis* and *E.*
926 *altidens* (Vekua, 1986, 1987; Hemmer et al., 2001; Tappen et al., 2002) and Dursunlu in
927 the generally more arid central Anatolia (1.0-0.78 Ma) is characterized by the high
928 frequency of *Equus caballus mosbachensis* and *E. altidens* (Güleç et al., 1999).

929 A later large faunal turnover is observed between the Epivillafranchian and the
930 Galerian faunas and is date dated to the Matuyama-Brunhes Boundary ca. 0.78 Ma. The
931 best record is derived from GBY. This dated to MOIS 18, is assigned to a wet and humid
932 stage apparent in the high frequency of cervids (*Dama* sp., *Cervus* cf. *Cervus elaphus* and
933 *Megalocerni* sp.) (Rabinovich et al., 2008). A small contribution of African taxa includes
934 *Bos buiaensis* (Martínez Navarro et al., 2009)

935 The sites of Denizli and Emirkaya-2 in Anatolia and Bear's cave, Tel Hesi and
936 'Ain Soda in the Levant are dated to the early part of the Middle Pleistocene roughly
937 between 0.5-0.43 Ma (Sen et al., 1991; Tchernov and Tsoukala, 1997; Dirks, 1998; Erten
938 et al., 2005). Taxa that appear in the region for the first time include *Panthera leo*, several
939 megacerines and *Capreolus* aff. *sussenborensis* while the continuation of archaic taxa
940 from earlier time periods include large taxa such as *Stephanorhinus hemitoechus* and *Bos*
941 *primigenius*. Unfortunately the taphonomic history of these sites precludes a detailed
942 paleoecological analysis. Bear's cave as its name implies is a carnivore den and presents
943 high frequencies of carnivore taxa and a minimal appearance of herbivore taxa, which
944 precludes an environmental reconstruction. The other sites have only scant remains.

945 Not surprisingly the assemblage of Yarimburgaz cave, situated in Thrace, the
946 European portion of modern Turkey, includes faunal elements from the Russian plains

such as *Lagurus transiens* and *Ochotona pusilla* which suggest the expansion of steppe environment in the southern Balkans, consistent with a cold and dry OIS 8 (von Koenigswald, 1998).

The sites of Qesem cave, Revadim and Holon, Hayonim cave layer E, Tabun cave layer E in the southern Levant are dated to the later part of the Middle Pleistocene between 0.43-0.13 Ma and provide a rich faunal assemblage for MOIS 8-6. The layers which dated to the earlier phase of the sequence (MOIS 7) are dominated by cervids most notably *Dama mesopotamica* whereas from MOIS 6, the increase in the mountain gazelle is noted. It has been hypothesized that the increasing frequencies of gazelle may indicate a climatically driven dispersal of an African arid species into the region (Stiner et al., 2004, 2009), which is supported by the presence of a single specimen of *Lycaon pictus* in Hayonim E (Stiner et al., 2001). However, isotopic analysis of teeth of both fallow deer and gazelles from the Mediterranean region from both Middle and Late Pleistocene sites of Qesem, Hayonim and Meged sites did not indicate any significant change in niche utilization between the ungulates, suggesting that the change in abundance may be due to human hunting preferences (Rowland, 2006). During this period there is no sound evidence for an additional dispersal from Africa following the Mid-Brunhes event of 0.43 Ma, represented by extinction of the larger fauna (Belmaker, 2009).

The modern terrestrial ecological signature of both the flora and the fauna of the region developed early in the Pleistocene. While the exact composition of species may have changed throughout the last 2.5 Million years, but the overall biome structure as a Mediterranean region surrounded by a more arid region in the South and East and

temperate region in the north did not alter significantly. The locations and extent of the borders between the sub-regions may have changed emphasizing the importance of understanding the role of ecotones in the region.

Discussion - faunal and hominins dispersals

Several trends can be observed in the paleoecological sequence of Western Asia during the Early and Middle Pleistocene. Within this long sequence, four major observations are recognized: 1. None of the environmental changes, and particularly in the Levant, were as dramatic as those recognized in higher latitudes (e.g., temperate Europe); 2. The climate changed gradually from more humid and closed habitats to more dry and open ones, through minor fluctuations generally correlated with the marine isotope data; 3. Punctuated faunal turnovers occurred in rough correlations with Western and Eastern Europe faunal units and 4. Hominin dispersals at the current state of research are not positively correlated with faunal turnovers (Table 2).

The evidence presented in this paper indicates that a few faunal dispersals between African and Eurasia occurred throughout the Late Pliocene through the Middle Pleistocene. There is a general agreement among scholars that these events were generally coincident with global climate changes. The main difficulty is attributing each dispersal event to the time when it took place. The reason is that the known sites and assemblages, described and discussed above, do not necessarily date these events. We tend to forget that the discovery of Lower Paleolithic sites is accidental and they cannot be considered as a continuous chronological sequence. Not surprisingly this is the nature of windows into the remote past, and one can argue that the dispersals of large mammals

occurred long before their bones were uncovered in the excavated sites. Relating the dispersal events to the known or conjectured age of an archaeological site may result in erroneous conclusions.

It should be stressed that the southwestern Asian fauna was established in this vast region since the Miocene and consisted of characteristic Palaearctic Eurasian element. The earliest dispersal from Africa of the generally woodland browser species (n=2) took place sometimes during the late Pliocene. It is recorded in the undated site of Bethlehem. The next migration was during the Early Pleistocene by several species marked by their adaptation to open grasslands (n=5). Their representation in the ‘Ubeidiya assemblages does not necessarily indicate the date of the first arrival of these taxa. A few additional browsers arrived at the time of the Jaramillo Event (n=2), and an additional grassland species at the Brunhes-Matuyama (n=1) as well as one taxon in the Mid-Brunhes (ca. 0.43 Ma; Belmaker, 2009).

The attempt to use the evidence from large mammals as a backdrop to understanding the tempo and mode of early hominin dispersals is generally unsuccessful. Contrary to what has been suggested (Martínez-Navarro et al, 2007, in press; Agustí et al., 2009a) it becomes apparent that hominin migrations into this region do not correlate with the arrivals of large mammals or specific climatic regimes. The earliest dated hominin dispersal is at the time of the Dmanisi (ca. 1.77 Ma) by bearers of a core and flake industry. The second, as no other Early Acheulian site was found and dated in southwestern Asia, is at ‘Ubeidiya (1.6-1.2 Ma).

During the Epivillafranchian, ca. 1.1 Ma, large mammalian fauna arrived in the Levant from Africa as attested by the fauna of Evron and Latamne. However, the lithic

1016 industry of these sites does not differ from other Early Acheulian assemblages, and in the
1017 lack of specific tool-types or new knapping techniques, we see no evidence for additional
1018 migration of African hominins into southwestern Asia.

1019 The site of GBY is interpreted to be the archaeological remains of a group of
1020 hominids that migrated from Africa (Bar-Yosef, 1987; Goren-Inbar and Saragusti, 1996).
1021 The lithic industry at GBY bears an African stamp and is still a rare occurrence in the
1022 Levant in spite of large surveys (e.g. Hours, 1981). The hominins had a definite
1023 preference for employing basalt as the main raw material although flint and limestone
1024 were present in the immediate vicinity. This choice, clearly expressed in high frequencies
1025 of cleavers, indicates the African origins of this group. It has been suggested (Bar-Yosef,
1026 1994, 1995) that this move was triggered by environmental change in East Africa that
1027 occurred around the Jaramillo subchron or the BMM. Interestingly, the makers of the
1028 GBY industry, either passed away after a time span of several dozens of millennia, or
1029 simply adopted the making of bifaces from flint with similar shapes to the common ones
1030 in the Levant.

1031 Human migrations occurred along the “Levantine Corridor” as defined by the
1032 paleontologists (e.g., Thomas, 1985). The Lower Paleolithic assemblages from el-
1033 Abassieh in Cairo (Bovier-Lapierre, 1926) may indicate that the Nile Valley was a
1034 possible segment of this route. The interior of the Arabian Peninsula was not an option
1035 due to the existence of the Saharan desert belt, since the end of the Miocene. Under
1036 interglacial conditions the northern penetration of the monsoonal system drastically
1037 changed the potentials for increasing amounts of resources in eastern Sahara (e.g.,

Neumann, 1989) and could have facilitated an alternative path for hominins and later Archaic *Homo sapiens* groups along the western shores of the Red Sea.

Thus, the absence of correlation between hominin and faunal dispersals is apparent and points to different underlying mechanisms for the two phenomenon. To reiterate the point we can discuss the situation in Arabia, which is beyond the scope of this paper. Regional surveys in the Arabian Peninsula have led to the identification of find spots and the collection of lithic assemblages with and without bifaces. Bifaces are reported solely from the western margins where they are made on a variety of raw materials such as flint, basalt, and metamorphic rocks. No bifaces have yet been found in the eastern Arabia that borders the Persian Gulf. Of special interest are the reports concerning sites or find spots along the Red Sea, another potential route of hominins. The excavation at Saffaqah (Whalen et al., 1983, 1984) provided a rich Acheulian assemblage made primarily of andesite with bifaces, cleavers, and numerous flakes. The depth of the deposits that contain artifacts (ca. 0.90m) probably reflects repeated occupations. Farther south in Yemen, excavations of open-air sites embedded in Pleistocene formations, many of which are rich in gravels or angular rock fragments, unearthed several series of core and flake assemblages and bifaces without animal bones (Amirkhanov, 1991). In addition, surface collections clearly indicate the presence of Late Acheulian industry. If no lithics remains are found in the anterior of Arabia during the Lower Paleolithic suggesting hominin migration into this region, then the site of An Nafud (Thomas et al., 1998) dated to the Early Pleistocene exhibits many of the African taxa present in the Southern Levant (*Crocota crocuta*; *Pelorovis* cf. *oldowayensis*; *Oryx* sp.) as well additional African alcelephines and bovids whereas Palaeartic species such as cervids

are notably absent. This would suggest that Africa fauna could disperse where early *Homo* could or would not. Further confirming the disjunct between the two dispersal patterns.

We should therefore return to the question concerning the nature of long distance migrations of hominins, whether it was based on “leapfrog” bursts of movements or gradual spreading out of small groups. Several scholars proposed that climatic changes determined the timing, mode and motivation for the out of Africa of early *Homo erectus/ergaster* (Potts, 1998; Dennell, 2004). It has been suggested that the expansion of grassland habitats into higher latitudes (30°N), during the Late Pliocene and Early Pleistocene climatic amelioration provided the favorite habitats that facilitated human survival (Vrba, 1988, 1995; Wynn, 2004). As the paleo-climatic information reflects alternating drier and wetter intervals it could have determined the closing and opening of natural corridors and thereby allowing *Homo* and other taxa to move from one region to the next (Vaks et al., 2002; Raia et al., 2006).

Indeed, the Aullan dispersal event (ca. 1.8 Ma) was associated with favorable ecological conditions permitting hominins and a few African taxa to reach Eurasia. However, the paleoecology of the Early Pleistocene sites across southwestern Asia (Dmanisi, ‘Ubeidiya, GBY) based on both flora and fauna indicates that the hominins survived in a mixed landscape of open parkland with forested surrounding hills (van Zeist and Bottema, 2009; Belmaker, 2009; Tappen, 2009; Belmaker, in press). Moreover, the prevailing environmental conditions are not sufficient to explain the differences between the lithic industries, whether ‘core and flake’ in the Caucasus region or the Early Acheulian in the Jordan Valley. Probably these two particular industries reflect the

1084 activities of different groups of hominins who by their traditional learning employed
1085 different stone artifacts.

1086 Successful adaptations to new ecological niches away from the savanna
1087 environment had its price. Probably more than one of the hominin lineages has
1088 disappeared i.e., simply died out, when they had too few members to keep a viable
1089 mating and reproductive system. Such local and temporal extinctions are possibly one of
1090 the reasons, together with low archaeological visibility, why the number of Lower
1091 Paleolithic sites, even in well-researched areas, is still so small. Therefore the current
1092 distribution of lithic industries across Eurasia is undoubtedly incomplete due to paucity of
1093 field research in several regions, indicating that there is no real cultural continuity and
1094 that what we view as a constant stream of migrants, was actually interrupted many times
1095 (e.g., Dennell, in press).

1096 Chronologically, the earliest hominins reaching southwestern Asia after leaving
1097 Africa were the makers of ‘core and flake’ industries, and is probably why this kind of
1098 simple way of obtaining sharp edges spread across Asia and Eastern Europe. It is not
1099 impossible that even migrants in later time (such as the makers of the Karari industry)
1100 carried the ‘core and flake’ production into the same areas. This would explain why the
1101 Acheulian contexts dated to 1.5 Ma to 0.25 Ma demonstrate interstratifications with those
1102 who manufactured cores and flakes (sometimes with additional types such as spheroids).

1103 The continuous occupation of southwestern Asia by the makers of the Acheulian
1104 industry needs to be stressed. The contrast with the neighboring regions is striking. No
1105 Acheulian sites are known from Iran, but plenty were recorded in Southern Asia (e.g.
1106 Petraglia, 1998). A few sites with bifaces are known in China beyond the “Movius line”

1107 (e.g., Hou et al, 2000). Between the Zagros and the western margins of the Indus valley
1108 there are suitable raw materials for making bifaces, but those who needed and knew how
1109 to make these tool-types were not present in these region. Natural boundaries such as the
1110 mountain ranges of the Caucasus limited early hominin moves as shown by the
1111 decreasing numbers of Acheulian handaxes (Liubin, 2002).

1112 In brief, we have no persuasive explanation why the Acheulian is not represented
1113 in the vast area between the Zagros and Baluchistan. A similar observation concerns
1114 Eastern Europe where only ‘core and flake’ industries were found and stand in marked
1115 difference with the proliferation of handaxes in Western Europe.

1116 While we agree that technological innovations and their social role are also
1117 considered as an important adaptation for dispersal (Carbonell et al., 1995; Larick and
1118 Ciochon, 1996), we wonder how exactly the technological innovations assisted in the
1119 survival or organization of those early hominin groups, and in particular given the lack of
1120 correlation between lithic technology, typology and environment. Detaching a few flakes
1121 from a nodule can be done in different ways, as it often depends of the fracture
1122 mechanics of the raw material, its size, shape and volume (Hovers and Braun, 2009 and
1123 papers therein).

1124 Therefore, if we accept the notion that adaptation to a specific environment during
1125 Lower Paleolithic times, did not influence the way stone tools were shaped, we have to
1126 adopt the position that it was due to the intrinsic behavioral capacities, unique to
1127 hominins, that facilitated the moves from Africa into new territories. It should be
1128 remembered that the initial dispersal of hominins has been attributed to both
1129 morphological and behavioral characteristics such as the capacity for long distance

1130 walking (Steudel, 1994), endurance running (Bramble and Lieberman, 2004), heat
1131 adaptation (Walker and Leakey, 1993), greater brain capacity (Aiello, 1993; Aiello and
1132 Wheeler, 1995) and social structure (Kroll, 1994). Even the release from tropical diseases
1133 that allowed for an increase in population size in higher latitudes (Bar Yosef and Belfer-
1134 Cohen, 2001), does not explain what happened to the various groups in Asia.

1135 The variability selection hypothesis states that the adaptability of hominins to a
1136 wide range of habitats and specifically to a variable climate in Africa may have provided
1137 a pre-adaptation to survival in novel environments (Potts, 1998, 2002). While we agree
1138 that the presence of hominins in a wide range of environments in Eurasia supports this
1139 hypothesis, however, this explanation is valid for the “longue durée” when a period of
1140 one million years or more is considered. In addition, hypothesizing what may constitute
1141 the pre-adaptability traits is rather intriguing. These may include an increase of resource
1142 exploitation by using of stone tools, wooden tools that did not survive, and fire. For the
1143 latter there is hardly any solid evidence prior to 0.8 Ma in GBY (Goren-Inbar et al., 2004;
1144 Alperson-Afil and Goren-Inbar, 2007; Alperson-Afil, 2008). Additional pre-adaptations
1145 would have been the importance of high-energy foods such as meat, tubers or other
1146 vegetal sources during the initial dispersal. The increase in body and brain size required a
1147 better quality diet and often the contribution of protein seems to have been the main
1148 source (Aiello and Wheeler, 1995; Cordain et al., 2000; Aiello et al., 2001; Cordain et al.,
1149 2001; Aiello and Wells, 2002). Hence, hominins evolved to become more active hunters
1150 and gatherers when compared to their ancestors, and these capacities required larger
1151 territories (Walker and Shipman, 1996). Specifically, a high proportion of meat in the diet
1152 was seen as a critical for the success of hominins and this proposition was supported,

1153 according to various authors, by the taphonomic analyses of faunal remains
1154 (Brantingham, 1998; Hemmer, 2000; Domínguez-Rodrigo et al., 2002; Domínguez-
1155 Rodrigo, 2002; Domínguez-Rodrigo and Pickering, 2003). The colonization of the
1156 temperate Europe after 0.7 Ma is thought as facilitated by a decrease in the number of
1157 carnivore competitors and the ability of hominins to acquire larger quantities of meat
1158 (Palombo and Mussi, 2006), although the role of large game hunting could signal more a
1159 social expression than the need for meat (J.D.Speth, personal communication).

1160 In sum, this paper provides an overview of the environmental conditions that
1161 prevailed during the Lower and Middle Pleistocene in southwestern Asia indicating that
1162 regional topographic configuration played a primary role in shaping the effects of
1163 climatic amplitude on the biotic responses of flora and fauna. The emphasis on climatic
1164 shifts and regional ecological variability is commonly seen as the sound background for
1165 the archaeological contexts and therefore facilitating the understanding hominin
1166 population dynamics. Hominins were able to disperse into regions that were beyond the
1167 their sub-tropical and tropical African homeland (Bosinski, 2006; Dennell 2009) and in
1168 routes, times and into environments unrelated to other large African taxa both carnivore
1169 and herbivore suggesting a unique biological, behavioral and cultural suite of characters
1170 which allowed them to do so. However, without a better understanding the reasons for
1171 success and failure of survival of various hominin migrant groups, and whether it
1172 depended on their ability to keep their mating and reproductive systems, we will need to
1173 resort to oversimplifications of the “human success”.

1174 1175 **Acknowledgements**

1176 We would like to thank the organizers of this volume for inviting us to participate.

1177 MB is funded by the American School of Prehistoric Research at Harvard University.

1178

1178

1179 **References**

1180 Abdul-Nayeem, M., 1990. Prehistory and Protohistory of the Arabian Peninsula. Volume I:

1181 Saudi Arabia. Hyderabad Publishers, Hyderabad.

1182 Agustí, J., Oms, O., Parés, J.M., Martínez-Navarro, B., Turq, A., 2000. Dating and correlation of

1183 early human occupation in the Baza Formation (Guadiz-Baza Basin, SE Spain). In:

1184 Lordkipanidze, D., Bar-Yosef, O., Otte, M. (Eds.), Early Humans at the Gates of Europe.

1185 ERAUL 92, Liège, pp. 113-122.

1186 Agustí, J., Blain, H.A., Cuenca-Bescos, G., Bailon, S., 2009a. Climate forcing of first hominid

1187 dispersal in Western Europe. *Journal of Human Evolution* 57, 815-821.

1188 Agustí, J., Vekua, A., Oms, O., Lordkipandize, D., Bukhsianidze, M., Kiladze, G., Rook, L.,

1189 2009b. The Plio-Pleistocene succession of Kvabebi (Georgia) and the background to the

1190 early human occupation of Southern Caucasus. *Quaternary Science Reviews* 28, 3275-

1191 3280.

1192 Aiello, L.C., 1993. Neocortex size, group size, and the evolution of language. *Current*

1193 *Anthropology* 34, 184-193.

1194 Aiello, L.C., Bates, N., Joffe, T., 2001. In defense of the expensive tissue hypothesis. In: Falk,

1195 D., Gibson, K.R. (Eds.), *Evolutionary Anatomy of the Primate Cerebral Cortex*.

1196 Cambridge University Press, Cambridge, pp. 57-78.

1197 Aiello, L.C., Wheeler, P., 1995. Expensive-tissue hypothesis: the brain and digestive system in

1198 human and primate evolution. *Current Anthropology* 36, 199-221.

1199 Aiello, L.C., Wells, J.C.K., 2002. Energetics and the evolution of the genus *Homo*. *Annual*

1200 *Review of Anthropology* 31, 323-338.

- 1201 Alperson-Afil, N., 2008. Continual fire-making by Hominins at Gesher-Benot Ya'akov, Israel.
1202 Quaternary Science Reviews 27, 1733-1739.
- 1203 Alperson-Afil, N., Goren - Inbar, N., 2007. Out of Africa and into Eurasia with controlled use of
1204 fire: Evidence from Gesher Benot Ya'akov, Israel. Archaeology, Ethnology and
1205 Anthropology of Eurasia 28, 63-78.
- 1206 Alperson-Afil, N., Richter, D., and Goren-Inbar, N. (2007). Phantom hearths and the use of Fire
1207 at Gesher Benot Ya'akov, Israel. PaleoAnthropology 7, 1-15.
- 1208 Amirkhanov, K.A., 1991. Paleolit Iuja Arabii. Nauka, Moscow.
- 1209 Antón, S.C., Swisher, C.C., 2004. Early dispersals of Homo from Africa. Annual Review of
1210 Anthropology 33, 271-296.
- 1211 Ariai, A., Thibault, C., 1975/77. Nouvelles précisions à propos de l'outillage paléolithique ancien
1212 sur galets du Khorassan (Iran). Paléorient 3, 101-108.
- 1213 Arribas, A., Palmqvist, P., 1999. On the ecological connection between saber-teeth and
1214 hominids: faunal dispersal events in the Lower Pleistocene and a review of the evidence
1215 of the first human arrival in Europe. Journal of Archaeological Science 26, 571-585.
- 1216 Ashkenazi, S., Klass, K., Mienis, H.K., Sprio, B., Abel, R., 2009. Fossil embryos and adult
1217 Viviparidae from the Early-Middle Pleistocene of Gesher Benot Y'akov, Israel: ecology,
1218 longevity and fecundity. Lethaia, 10.1111/j.1502-3931.2009.00178.x.
- 1219 Bar-Yosef, O., 1987. Pleistocene connections between Africa and Southwest Asia: An
1220 archaeological perspective. The African Archaeological Review 5, 29-38.
- 1221 Bar-Yosef, O., 1994. The Lower Paleolithic of the Near East. Journal of World Prehistory 8,
1222 211-265.
- 1223 Bar-Yosef, O., 1995. The Origins of Modern Humans. In: Levy, T.E. (Ed.), Archaeology of

- 1224 Society in the Holy Land. Facts on File, New York, pp. 110-123.
- 1225 Bar-Yosef, O., 1998a. Early colonizations and cultural continuities in the Lower Palaeolithic of
 1226 Western Asia. In: Petraglia, M.D., Korisettar, R. (Eds.), Early Human Behaviour in
 1227 Global Context: The Rise and diversity of the Lower Palaeolithic record. Routledge,
 1228 London, pp. 221-279.
- 1229 Bar-Yosef, O., 1998b. Jordan Prehistory: A View from the West. In: Henry, D.O. (Ed.), The
 1230 Prehistoric Archaeology of Jordan. Archaeopress, Oxford, pp. 162-178.
- 1231 Bar-Yosef, O., Tchernov, E., 1972. On the Palaeo-Ecological History of the Site of 'Ubeidiya.
 1232 Israel Academy of Sciences and Humanities, Jerusalem.
- 1233 Bar-Yosef, O., Goren-Inbar, N., 1993. The Lithic Assemblages of 'Ubeidiya, a Lower Paleolithic
 1234 Site in the Jordan Valley. The Institute of Archaeology, The Hebrew University of
 1235 Jerusalem, Jerusalem.
- 1236 Bar-Yosef, O., Belfer-Cohen, A., 2001. From Africa to Eurasia - early dispersals. Quaternary
 1237 International 75, 19-28.
- 1238 Barkai, R., Gopher, A., Lauritzen, S.E., Frumkin, A., 2003. Uranium Series dates from Qesem
 1239 Cave, Israel, and the end of the Lower Paleolithic. Nature 423, 977-979.
- 1240 Barkai, R., Gopher, A., and Shimelmitz, R. 2006. Middle Pleistocene Blade Production in the
 1241 Levant: An Amudian Assemblage from Qesem Cave, Israel. Eurasian Prehistory 3, 39-
 1242 74.
- 1243 Barzilai, O., Buller, A. M., and Ackermann, O. (2006). Kefar Menachem West: A Lower
 1244 Paleolithic site in the Southern, Shephela, Israel. Journal of The Israel Prehistoric Society
 1245 36, 7-38.
- 1246 Behrensmeier, A.K., 2006. Climate Change and Human Evolution. Science 311, 476-478.

- 1247 Belmaker, M., 2006. Community structure through time: 'Ubeidiya, a Lower Pleistocene site as a
1248 case study. Ph.D. Dissertation, The Hebrew University of Jerusalem.
- 1249 Belmaker, M., 2009. Hominin adaptability and patterns of faunal turnover in the Lower-Middle
1250 Pleistocene transition in the Levant. In: Camps, M., Chauhan, P.R. (Eds.), A sourcebook
1251 of Paleolithic transitions: Methods, Theories and Interpretations. Springer, pp. 211-227.
- 1252 Belmaker, M., 2010. The presence of a large cercopithecine (cf. *Theropithecus* sp.) in the
1253 'Ubeidiya Formation (Early Pleistocene 'Ubeidiya). *Journal of Human Evolution* 58,79-
1254 89.
- 1255 Belmaker, M., in press. Early Pleistocene faunal connections between Africa and Eurasia: An
1256 ecological perspective. In: Fleagle, J., Grine, F., Leakey, R., J.J, S. (Eds.), *Out of Africa*
1257 *I: Who, Where and When*. Springer. New York.
- 1258 Belmaker, M., Tchernov, E., Condemi, S., Bar-Yosef, J., 2002. New evidence for hominid
1259 presence in the Lower Pleistocene in the Southern Levant. *Journal of Human Evolution*
1260 43, 43-56.
- 1261 Bennett, K.D., 1997. *Evolution and Ecology: The Pace of Life*. Cambridge University Press,
1262 Cambridge.
- 1263 Braidwood, R., Howe, B., (Eds.) 1960. *Prehistoric investigations in Iraqi Kurdistan*. Oriental
1264 Institute, Chicago.
- 1265 Bosinski, G., 2006. Les premiers peuplements de l'Europe centrale et de l'Est. *C.R. Palevol* 5,
1266 311-317.
- 1267 Bovier-Lapierre, R.P., 1926. Les gisements paléolithiques de la plaine de l'Abbassieh. *Bulletin*
1268 *de l'Institut d'Égypte* b, 257-275.
- 1269 Bramble, D.M., Lieberman, D.E., 2004. Endurance running and the evolution of *Homo*. *Nature*

- 1270 432, 345-352.
- 1271 Brantingham, J.P., 1998. Hominid-carnivore coevolution and invasion of the predatory guild.
1272 *Journal of Anthropological Archaeology* 17, 327-353.
- 1273 Braun, D., Ron, H., Marco, S., 1991. Magnetostratigraphy of the hominid tool-bearing Erk el
1274 Ahmar formation in the northern Dead Sea Rift. *Israel Journal of Earth Sciences* 40, 191-
1275 197.
- 1276 Braüer, G., Schultz, M., 1996. The morphological affinities of the Plio-Pleistocene mandible
1277 from Dmanisi, Georgia. *Journal of Human Evolution* 30, 445-481.
- 1278 Butzer, K., 1958. Quaternary Stratigraphy and Climate in the Near East. *Bonner Geographische*
1279 *Abhandlungen*, Fred Dummlers Verlag,
- 1280 Carbonell, E., Bermudez de Castro, J.M., Arsuaga, J.L., Diez, J.C., Rosas, A., Cuenca-Bescos,
1281 G., Sala, R., Mosquera, M., Rodriguez, X.P., 1995. Lower Pleistocene Hominids and
1282 Artifacts from Atapuerca-TD6 (Spain). *Science* 269, 826-830.
- 1283 Caloi, L., Palombo, M.R., 1997. Biochronology of large mammals in the early and middle
1284 Pleistocene on the Italian peninsula. *Hystrix* 9, 3-12.
- 1285 Chazan, M., Kolska Horwitz, L., (Eds.) 2007. Holon. A Lower Paleolithic Site in Israel. Peabody
1286 Museum of Archaeology and Ethnology, Cambridge MA.
- 1287 Cerling, T.E., Hay, R.L., 1986. An isotopic study of paleosol carbonates from Olduvai Gorge.
1288 *Quaternary Research* 25, 63-78.
- 1289 Clark, J.D., 1961. Fractured Chert Specimens from the Lower Pleistocene Bethlehem Beds,
1290 Israel. *Bulletin of the British Museum* 5, 73-99.
- 1291 Clark, J.D., 1967. The middle Acheulian site at Latamne, northern Syria. *Quaternaria* 9, 1-68.
- 1292 Copeland, L., Hours, F., 1981. La fin de l'Acheuléen et l'avènement du Paléolithique Moyen en

1293 Syrie. In: Cauvin, J., Sanlaville, P. (Eds.), La Préhistoire du Levant. CNRS, Paris, pp.
 1294 225-238.
 1295 Copeland, L., Hours, F., 1983. Le Yabroudien d'El Kowm (Syrie) et sa place dans le
 1296 Paléolithique du Levant. *Paléorient* 9, 21-37.
 1297 Copeland, L., Hours, F., 1989. The Hammer on the Rock: Studies in the Early
 1298 Palaeolithic of Azraq, Jordan. BAR, Oxford.
 1299 Cordain, L., Miller, J.B., Eaton, S.B., Mann, N., Holt, S.H.A., Speth, J.D., 2000. Plant-
 1300 animal subsistence ratios and macronutrient energy estimations in worldwide
 1301 hunter-gatherer diets. *American Journal of Clinical Nutrition* 71, 682-692.
 1302 Cordain, L., Watkins, B.A., Mann, N.J., 2001. Fatty acid composition and energy density of
 1303 foods available to African hominids. Evolutionary implications for human brain
 1304 development. In: Simopoulos, A.P., Pavlou, K.N. (Eds.), *Nutrition and Fitness: Metabolic*
 1305 *Studies in Health and Disease*. Karger, Basel, pp. 114-161.
 1306 Danin, A., 1988. Flora and vegetation of Israel and adjacent areas. In: Yom-Tov, Y., Tchernov,
 1307 E. (Eds.), *The Zoogeography of Israel. The Distribution and Abundance at a*
 1308 *Zoogeographical Crossroad*. Dr. W. Junk Publishers, Dordrecht, pp.129-158.
 1309 de Heinzelin, J., 1966. Geological observations near Latamne. *Annales Archéologiques Arabes*
 1310 *Syriennes* 16, 115-120.
 1311 Dennell, R.W., 2004. Hominid dispersals and Asian biogeography during the Lower and Early
 1312 Middle Pleistocene, c. 2.0 - 0.5 Mya. *Asian Perspectives* 43, 205-226.
 1313 de Lumley, H., Nioradze, M., Barsky, D., Cauche, D., Celiberti, V., Nioradze, G., Notter, O., D.,
 1314 Z., Lordkipanidze, D., 2005. les industries lithiques preoldowayennes du debut du
 1315 Paléolithique inferieur du site de Dmanisi en Georgie. *L'Anthropologie* 109, 1-182.

- 1316 Dennell, R.W., 2004. Hominid dispersals and Asian biogeography during the Lower and Early
1317 Middle Pleistocene, c. 2.0 - 0.5 Mya. *Asian Perspectives* 43, 205-226.
- 1318 Dennell, R.W., in press. "Out of Africa 1" and the evidence for hominins in Eurasia before 0.5
1319 Ma: problems and prospects. In: Fleagle, J., Grine, F., Leakey, R., Shea, J.J. (Eds.), *Out*
1320 *of Africa I: Who, Where and When*. Springer, New York.
- 1321 D'Errico, F., Nowell, A., Bar-Yosef, O., Zilhao, J., Mithen, S., Wynn, T., Marshack, A., Close,
1322 A.E., 2000. A new look at the Berekhat Ram figurine: Implications for the origins of
1323 symbolism. *Cambridge Archaeological Journal* 10, 123-167.
- 1324 Derricourt, R., 2006. Getting "Out of Africa": Sea crossing, land crossings and culture in the
1325 hominin migrations. *Journal of World Prehistory* 19, 119-132.
- 1326 Dirks, W., Schunrenberger, D.W., 1998. Preliminary account of Pleistocene mammals from 'Ain
1327 Soda, Azraq Basin, Jordan. *Journal of Vertebrate Paleontology* 18, 38A-39A.
- 1328 Dodonov, A.E., Deviatkin, E.V., Ranov, V.A., 1993. Latamne formation in the Orontes river
1329 valley. In: Sanlaville, P., Besançon, J., Copeland, L., Muhesen, S. (Eds.), *Le*
1330 *Paléolithique de la Vallée Moyenne de l'Oronte (Syrie)*. BAR International Series 587,
1331 Oxford, pp. 189-194.
- 1332 Domínguez-Rodrigo, M., 2002. Hunting and scavenging by early humans: the state of the debate.
1333 *Journal of World Prehistory* 16, 1-54.
- 1334 Domínguez-Rodrigo, M., De la Torre, I., De Luque, L., Alcalá, L., Mora, R., Serrallonga, J.,
1335 Medina, V., 2002. The ST Site Complex at Peninj, West Lake Natron, Tanzania:
1336 implications for early hominid behavioural models. *Journal of Archaeological Science*
1337 29, 639-665.
- 1338 Domínguez-Rodrigo, M., Pickering, T.R., 2003. Early Hominid Hunting and Scavenging: A

- 1339 Zooarcheological Review. *Evolutionary Anthropology* 12, 275-282.
- 1340 Dzaparidze, V., Bosinski, G., Bugianisvili, T., Gabunia, L., Justus, A., Klopotovskaja, N.,
 1341 Kvavadze, E., Lordkipanidze, D., Majsuradze, G., Mgeladze, N., Nioradze, M.,
 1342 Pavlenisvili, E., Schminke, H.-U., Sologasvili, D., Tusabramsvili, Tvalcrelidze, M.,
 1343 Vekua, A., 1992. Der altpaläolithische Fundplatz Dmanisi in Georgen (Kaukasus).
 1344 *Jahrbuch Römisch-Germanischen Zentralmuseums Mainz* 36, 67-116.
- 1345 Enzel, Y., Amit, R., Dayan, U., Crouvi, O., Kahana, R., Ziv, B., Sharon, D., 2008. The climatic
 1346 and physiographic controls of the eastern Mediterranean over the late Pleistocene
 1347 climates in the southern Levant and its neighboring deserts. *Global and Planetary Change*
 1348 60, 165-192.
- 1349 Erten, H., Sen, S., Özkul, M., 2005. Pleistocene mammals from travertine deposits of the Denizli
 1350 basin (SW Turkey). *Annales de paléontologie* 91, 267-278.
- 1351 Feibel, C.S., 2004. Quaternary lake margins of the Levant Rift Valley. In: Speth, J.D., Goren-
 1352 Inbar, N. (Eds.), *Human Paleoecology in the Levantine Corridor*. Oxbow Publishers,
 1353 Oxford, pp. 21-36.
- 1354 Frumkin, A., Stein, M., 2004. The Sahara–East Mediterranean dust and climate connection
 1355 revealed by strontium and uranium isotopes in a Jerusalem speleothem. *Earth and*
 1356 *Planetary Science Letters* 217, 451-464.
- 1357 Gabunia, L., Vekua, A., 1990. L'Évolution du paléoenvironnement au cours de l'Anthropogène
 1358 en Géorgie (Transcaucasie). *L'Anthropologie* 94, 643-650.
- 1359 Gabunia, L., Vekua, A., 1995. A Plio-Pleistocene hominid from Dmanisi, East Georgia,
 1360 Caucasus. *Nature* 373, 509-512.
- 1361 Gabunia, L., Vekua, A., Lordkipanidze, D., Swisher, C.C., Ferring, R., Justus, A., Nioradze, M.,

- 1362 Tvalchrelidze, M., Antón, S.C., Bosinski, G., Jöris, O., de Lumley, M.-A., Majsuradze,
1363 G., Mouskhelishvili, A., 2000. Earliest Pleistocene hominid cranial remains from
1364 Dmanisi, Republic of Georgia: taxonomy, geological setting, and age. *Science* 288, 1019-
1365 1025.
- 1366 Gabunia, L., de Lumley, M.-A., Vekua, A., Lordkipanidze, D., de Lumley, H., 2002. Découverte
1367 d'un nouvel hominidé à Dmanisi (Transcaucasie, Géorgie). *C. R. Palevol* 1, 243-253.
- 1368 Gardner, E.W., Bate, D.M.A., 1937. The bone-bearing beds of Bethlehem: their fauna and
1369 industry. *Nature* 140, 43-433.
- 1370 Geraads, D., and Tchernov, E. (1983). Fémurs humaine de Pléistocène moyen de Gesher Benot
1371 Ya'aqov (Israël). *L'Anthropologie* 87, 138-141.
- 1372 Gilead, D., 1970. Handaxe industries in Israel and the Near East. *World Archaeology* 2, 1-11.
- 1373 Gilead, D., Israel, M., 1975. An Early Palaeolithic site at Kefar Menahem: Preliminary report.
1374 *Tel Aviv* 2, 1-12.
- 1375 Ginat, H., Zilberman, E., Saragusti, I., 2003. Early Pleistocene lake deposits and Lower
1376 Paleolithic finds in Nahal (wadi) Zihor, Southern Negev desert, Israel. *Quaternary*
1377 *Research* 59, 445-458.
- 1378 Gisis, I., Bar-Yosef, O., 1974. New Excavation in Zuttiyeh Cave, Wadi Amud, Israel. *Paleorient*
1379 2, 175-180.
- 1380 Grün, R., Schwarcz, H.P., Stringer, C.B., 1991. ESR dating of teeth from Garrod's Tabun cave
1381 collection. *Journal of Human Evolution* 20, 231-248.
- 1382 Grün, R., Stringer, C., 2000. Tabun revisited: revised ESR chronology and new ESR and U-
1383 series analyses of dental material from Tabun C1. *Journal of Human Evolution* 39, 601-
1384 612.
- 1385 Gopher, A., Barkai, R., Shimelmitz, R., Khalaily, M., Lemorini, C., Hershkovitz, I., and Stiner,

- 1386 M. (2005). Qesem Cave: An Amudian Site in Central Israel. In "Journal of the Israel
1387 Prehistoric Society." (M. Haeven, Ed.), pp. 69-92. The Israel Prehistoric Society,
1388 Jerusalem.
- 1389 Goren, N., 1979. An Upper Acheulian industry from the Golan Heights. *Quartär* 29,30, 105-121.
- 1390 Goren-Inbar, N., 1985. The lithic assemblage of the Berekhat Ram Acheulian Site, Golan
1391 Heights. *Paléorient* 11, 7-28.
- 1392 Goren-Inbar, N. (1986). A figurine from the Acheulian site of Berekhat Ram. *Journal of the*
1393 *Israel Prehistoric Society* 19, 7-12.
- 1394 Goren-Inbar, N., 1995. The Lower Paleolithic of Israel. In: Levy, T. (Ed.), *The Archaeology of*
1395 *Society in the Holy Land*. Leicester University Press, London, pp. 93-109.
- 1396 Goren-Inbar, N., Zohar, I., Ben-Ami, D., 1991. A new look at old cleavers--Gesher Benot
1397 Ya'aqov. *Journal of the Israel Prehistoric Society* 24, 7-33.
- 1398 Goren-Inbar, N., Belitzky, S., Goren, Y., Rabinovich, R., Saragusti, I., 1992a. Gesher Benot
1399 Ya'aqov – the 'Bar': an Acheulian assemblage. *Geoarchaeology* 7, 27-40.
- 1400 Goren-Inbar, N., Belitzky, S., Verosub, K., Werker, E., Kislev, M., Heimann, A., Carmi, I.,
1401 Rosenfeld, A., 1992b. New discoveries at the Middle Pleistocene Acheulean site of
1402 Gesher Benot Ya'aqov, Israel. *Quaternary Research* 38, 117-128.
- 1403 Goren - Inbar, N., Saragusti, I., 1996. An Acheulian biface assemblage from Gesher Benot
1404 Ya'aqov, Israel: Indications of African Affinities. *Journal of Field Archaeology* 23, 15-
1405 30.
- 1406 Goren - Inbar, N., Feibel, C.S., Verosub, K.L., Melamed, Y., Kislev, M.E., Tchernov, E.,
1407 Saragusti, I., 2000. Pleistocene milestones on the out of Africa corridor at Gesher Benot
1408 Yaaqov, Israel. *Science* 289, 944-947.

- 1409 Goren-Inbar, N., Alpers, N., Kislev, M.E., Simchoni, O., Melamed, Y., Ben-Nun, A., Werker,
1410 E., 2004. Evidence of hominin control of fire at Gesher Benot Ya'akov, Israel. *Science*
1411 304, 725-727.
- 1412 Guérin, C., Eisenmann, V., Faure, M., 1993. Les grands mammifères du gisement pléistocène
1413 moyen de Latamné (vallée de l'Oronte, Syrie). In: Sanlaville, P., Besançon, J., Copeland,
1414 L., Muhesen, S. (Eds.), *Le Paléolithique de la vallée moyenne de l'Oronte (Syrie)*. BAR
1415 International Series 587, Oxford, pp. 169-178.
- 1416 Güleç, E., Howell, F.C., White, T.D., 1999. Dursunlu - A new Lower Pleistocene faunal and
1417 artifact-bearing locality in southern Anatolia. In: Ullrich, H. (Ed.), *Hominid Evolution:
1418 Lifestyles and Survival Strategies*. Edition Archaea, Berlin; Gelsenkirchen, Germany, pp.
1419 349-364.
- 1420 Gvirtzman, G., Wieder, M., Marder, O., Khalaily, H., Rabinovich, R., Ron, H., 1999. Geological
1421 and pedological aspects of an Early Paleolithic site: Revadim, Central Coastal Plain,
1422 Israel. *Geoarchaeology* 14, 101-126.
- 1423 Harrison, D.L., 1972. *Mammals of Arabia*. Ernst Benn, London.
- 1424 Harrison, D.L., Bates, P.J.J., 1991. *The Mammals of Arabia*. Harrison Zoological Museum,
1425 Sevenoaks, Kent.
- 1426 Hemmer, H., 2000. Out of Asia: a paleoecological scenario of man and his carnivorous
1427 competitors in the European Lower Pleistocene. In: Lordkipanidze, D., Bar-Yosef, O.,
1428 Otte, M. (Eds.), *Early Humans at the Gates of Europe. Etudes et Recherches*
1429 *Archéologiques de l'Université de Liège*, Liège, pp. 99-106.
- 1430 Hemmer, H., Kahlke, R.-D., Vekua, A.K., 2001. The jaguar *Panthera onca gombaszegensis*
1431 (Kretzoi, 1938) (Carnivora: Felidae) in the late Lower Pleistocene of Akhalkalaki (South

- 1432 Georgia; Transcaucasus) and its evolutionary and ecological significant. *Geobios* 34,
1433 475-486.
- 1434 Hooijer, D.A., 1958. An early Pleistocene mammalian fauna from Bethlehem. *Bulletin of the*
1435 *British Museum of Natural History, London, Geology* 3, 265-292.
- 1436 Hooijer, D.A., 1962. Middle Pleistocene mammals from Latamne, Orontes valley. *Annales*
1437 *Archéologiques de Syrie* 11, 117-132.
- 1438 Horowitz, A., 1979. *The Quaternary of Israel*. Academic Press, New York.
- 1439 Horowitz, A., 1988. The quaternary environments and paleogeography in Israel. In: Yom-Tov,
1440 Y., Tchernov, E. (Eds.), *The Zoogeography of Israel*. Dr. W.Junks Publishers, Dordrecht,
1441 Netherlands, pp. 35-57.
- 1442 Horowitz, A., 2001. *The Jordan Rift Valley*. A. A. Balkema, Rotterdam.
- 1443 Hou, Y., Potts, R., Yuan, B., Guo, Z., Deino, A., Wang, W., Clark, J., Xie, G., and Huang, W.
1444 (2000). Mid-Pleistocene Acheulean-like Stone Technology of the Bose Basin, South
1445 China. *Science* 287, 1622–1626.
- 1446 Hours, F., 1975. The Lower Paleolithic of Lebanon and Syria. In: Wendorf, E., Marks, A.E.
1447 (Eds.), *Problems in Prehistory: North Africa and the Levant*. S.M.U. Press, Dallas, pp.
1448 249-271.
- 1449 Hours, F., 1981. Le Paléolithique inférieur de la Syrie et du Liban. Le point de la question en
1450 1980. In: Sanlaville, P., Cauvin, J. (Eds.), *Préhistoire du Levant*. Maison de l'Orient,
1451 Lyon, pp. 165-184.
- 1452 Hours, F., Le Tensorer, J.M., Muhesen, S., Yalçinkaya, I., 1983. Premiers travaux sur le site de
1453 Nadaouyeh (El-Kowm, Syrie). *Paléorient* 9, 5-14.
- 1454 Hovers, E., Braun, D.R., 2009. *Interdisciplinary Approaches to the Oldowan*. Springer.

1455 Dordrecht.

1456 Hume, G.W., 1976. The Ladizian: An Industry of the Asian Chopper-Chopping Tool Complex in
 1457 Iranian Baluchistan. Dorrance and Company, Philadelphia.

1458 Issar, A., 1979. Stratigraphy and paleoclimates of the Pleistocene of central and northern Israel.
 1459 Palaeogeography, Palaeoclimatology, Palaeoecology 29, 262-280.

1460 Jacobs, B.F., Kingston, J.D., L.L., J., 1999. The origin of grass-dominated ecosystems. Annals of
 1461 the Missouri Botanical Garden 86, 590-643.

1462 Jelinek, A., 1981. The Middle Paleolithic of the Southern Levant from the Perspective of Tabun
 1463 Cave. In: Cauvin, J., Sanlaville, P. (Eds.), *Préhistoire du Levant*. Paris: Editions du
 1464 CNRS, pp. 265-280.

1465 Jelinek, A., 1982a. The Middle Paleolithic in the Southern Levant from the Perspective of the
 1466 Tabun Cave. In: Ronen, A. (Ed.), *The Transition from Lower to Middle Paleolithic and
 1467 the Origins of Modern Man*. British Archaeological Reports, Oxford, pp. 57-104.

1468 Jelinek, A., 1982b. The Middle Paleolithic in the Southern Levant with comments on the
 1469 appearance of modern *Homo sapiens*. In: Ronen, A. (Ed.), *The Transition from Lower to
 1470 Middle Paleolithic and the Origins of Modern Man*. British Archaeological Reports,
 1471 Oxford, pp. 57-104.

1472 Jöris, O. (2008). "Der Altpaläolithische Fundplatz Dmanisi, (Georgien, Kaukasus)." *Römisch-
 1473 Germanisches Zentralmuseum, Mainz*.

1474 Karkanas, P., Shahack-Gross, R., Ayalon, A., Bar-Matthews, M., Barkai, R., Frumkin, A.,
 1475 Gopher, A., and Stiner, M. C. (2007). Evidence for habitual use of fire at the end of the
 1476 Lower Paleolithic: Site-formation processes at Qesem Cave, Israel. *Journal of Human
 1477 Evolution* 53, 197-212.

- 1478 Kikodze, Z.K., 1986. Bifas-kolon v asele Kavkaza. Vestnik Gos. Muzeja Gruzii 38, 55-69.
- 1479 Kingston, J.D., 2007. Shifting adaptive landscapes: progress and challenges in reconstructing
1480 early hominid environments. Yearbook of Physical Anthropology 50, 20-58.
- 1481 Kolska - Horwitz, L., Tchernov, E., 1989. The Late Acheulian Fauna from Oumm Zinat. Journal
1482 of the Israel Prehistoric Society 22, 7*-14*.
- 1483 Kroll, E.M., 1994. Behavioral implications of Plio-Pleistocene archaeological site structure.
1484 Journal of Human Evolution 27, 1-3.
- 1485 Larick, R., Ciochon, R.L., 1996. The African emergence and early Asian dispersal of the genus
1486 Homo. American Scientist 84, 538-551.
- 1487 Laukhin, S.A., Ronen, A., Pospelova, G.A., Sharonova, Z.V., Ranov, V.A., Burdukiewicz, J.-M.,
1488 Volgina, V.A., Tsatskin, A., 2001. New data on the geology and Geochronology of the
1489 Lower Palaeolithic site Bizat Ruhama in the Southern Levant. Paléorient 27, 69-80.
- 1490 Lemorini, C., Stiner, M., Gopher, A., R., S., Barkai, R., 2006. Use-wear analysis of an Amudian
1491 laminar assemblage from the Acheuleo-Yabrudian of Qesem Cave, Israel. Journal of
1492 Archaeological Science 33, 921-934.
- 1493 Le Tensorer, J. L. (2004). Nouvelles fouilles à Hummal (El Kown, Syrie centrale) premiers
1494 resultants (1997-2001). In "From the River to the Sea: The Palaeolithic and the Neolithic
1495 on the Euphrates and in the Northern Levant." (O. Aurenche, M. L. Mièrè, and P.
1496 Sanlaville, Eds.), Archaeopress. pp. 223-240.
- 1497 Le Tensorer, J.-M., Muhesen, S., Jagher, R., 1993. Nadaouiyeh I Aïn Askar: Une grande
1498 séquence paléolithique du bassin d'El Kowm (Syrie), premiers résultats, fouilles 1989-
1499 1992. In: Cauvin, J. (Ed.), Cahiers de l'Euphrate 7. Editions Recherche sur les
1500 Civilisations, Paris, pp. 11-36.

- 1501 Liubin, V. P. (2002). "L'Acheuléen du Caucase." ERAUL, Liège.
- 1502 Liubin, V.P., Bosinski, G., 1995. The earliest occupation of the Caucasus region. In: Roebroeks,
1503 W., van Kolfschoten, T. (Eds.), The Earliest Occupation of Europe. University of Leiden,
1504 Leiden, pp. 207-253.
- 1505 Lorch, Y., 1966. A Pleistocene florule from the Jordan Valley. Israel Journal of Botany 15, 31-
1506 34.
- 1507 Lordkipanidze, D., Jashashvili, T., Vekua, A., Ponce de Leon, M., Zolokofer, C.P.E., Rightmire,
1508 G.P., Pontzer, H., Ferring, R., Oms, O., Tappen, M., Bushsianidze, M., Agusti, J.,
1509 Kahlke, R.D., Kildze, G., Martinez Navaro, B., Mouskhelishvili, A., Nioradze, M., Rook,
1510 L., 2007. Postcranial evidence from early *Homo* from Dmanisi, Georgia. Nature 449,
1511 305-310.
- 1512 Marder, O., Gvartzman, G., Ron, H., Khalaily, H., Wieder, M., Bankirer, R., Rabinovitz, R.,
1513 Porat, N., Saragusti, I., 1999. The Lower Palaeolithic site of Revadim Quarry,
1514 preliminary finds. Journal of the Israel Prehistoric Society - Mitekufat Haeven 28, 21-53.
- 1515 Marder, O., Milevski, I., Matskevich, Z., 2006. The handaxes of Revadim Quarry: typo-
1516 technological considerations and aspects of intra-site variability. In: Goren-Inbar, N.,
1517 Sharon, G. (Eds.), Axe Age. Acheulian Toolmaking from Quarry to Discard. Equinox,
1518 London, pp. 223-242.
- 1519 Martínez-Navarro, B., 2004. Hippos, pigs, bovids, saber-toothed tigers, monkeys, and hominids:
1520 dispersals through the Levantine corridor during late Pliocene and early Pleistocene
1521 times. In: Goren-Inbar, N., Speth, J.D. (Eds.), Human Paleoecology in the Levantine
1522 Corridor. Oxbow Books, Oxford, pp. 37-52.
- 1523 Martínez-Navarro, B., Palmqvist, P., 1995. Presence of the African Machairdont Megantereon

1524 whitei (Broom, 1937) (Felidae, Carnivora, Mammalia) in the Lower Pleistocene site of
1525 Venta Micene (Orce, Granada, Spain), with some considerations on the origin, evolution
1526 and dispersal of the genus. *Journal of Archaeological Science* 22, 569-582.

1527 Martínez - Navarro, B., Pérez-Carlos, J.-A., Palombo, M.R., Rook, L., Palmqvist, P., 2007. The
1528 Olduvai buffalo *Pelorovis* and the origin of *Bos*. *Quaternary Research* 68, 220-226.

1529 Martínez-Navarro, B., Belmaker, M., Bar-Yosef, O., 2009. The large carnivores from 'Ubeidiya
1530 (early Pleistocene, Israel): biochronological and biogeographical implications. *Journal of*
1531 *Human Evolution* 56, 514-524.

1532 Martínez-Navarro, B., Rook, L., Papini, M., Libsekal, Y., 2009. A new species of bull from the
1533 Early Pleistocene paleoanthropological site of Buia (Eritrea): Parallelism on the dispersal
1534 of the genus *Bos* and the Acheulian culture. *Quaternary International* (in press).

1535 Marshack, A., 1997. The Berekhat Ram figurine: a Late Acheulian carving from the Middle
1536 East. *Antiquity* 72, 327-337.

1537 McCown, T.D., Keith, A., 1939. *The Stone Age of Mt. Carmel, Volume 2: The Fossil Human*
1538 *Remains from the Levallois-Mousterian*. Oxford University Press, Oxford.

1539 Mercier, N., Valladas, H., Valladas, G., Reyss, J.L., Jelinek, A., Meignen, L., Joron, J.L., 1995.
1540 TL dates of burnt flints from Jelinek's excavations at Tabun and their implications.
1541 *Journal of Archaeological Science* 22, 495-510.

1542 Mercier, N., Valladas, H., Froget, L., Joron, J.L., Ronen, A., 2000. Datation par la
1543 thermoluminescence de la base du gisement paléolithique de Tabun (Mont Carmel,
1544 Israël). *C. R. Acad. Sci.* 330, 731-738.

1545 Minzoni-Déroche, A., Sanlaville, P., 1988. Le Paléolithique inférieur de la région de Gaziantep.
1546 *Paléorient* 14, 87-98.

- 1547 Muhesen, S., 1985. L'Acheuléen récent évoulé de Syrie. (Ed.), BAR International Series 248,
1548 Oxford, pp.
- 1549 Muhesen, S., 1993. L'Acheuléen récent évolué de l'Oronte. In: Sanlaville, P., Besançon, J.,
1550 Copeland, L., Muhesen, L. (Eds.), Le Paléolithique de la vallée moyenne de l'Oronte.
1551 BAR International Series 587, Oxford, pp. 145-166.
- 1552 Neumann, K., 1989. Holocene Vegetation of the Eastern Sahara: Charcoal from Prehistoric Sites.
1553 African Archaeological Review 7, 97-116.
- 1554 Neuville, R., 1951. Le Paléolithique et le Mésolithique de Désert de Judée. Masson et Cie,
1555 Editeurs, Paris.
- 1556 Ohel, M., 1991. Prehistoric survey of the Baram Plateau. Palestine Exploration Quarterly Jan-
1557 June, 32-47.
- 1558 Opdyke, N.D., Lindsay, E., Kukla, G., 1983. Evidence for earlier date of 'Ubeidiya, Israel
1559 hominid site. Nature 304, 375.
- 1560 Palombo, M.R., Mussi, M., 2006. Large mammals guild at the time of the first human
1561 colonization of Europe: The case of the Italian Pleistocene record. Quaternary
1562 International 149, 94-103.
- 1563 Petraglia, M. D., 1998. The Lower Palaeolithic of India and its bearing on the Asian record. In
1564 "Early Human Behaviour in Global Context: the Rise and Diversity of the Lower
1565 Palaeolithic Record." (M. D. Petraglia, and R. Korisettar, Eds.), pp. 343-390. Routledge,
1566 London.
- 1567 Porat, N., Chazan, M., Schwarcz, H., Horwitz, L.K., 2002. Timing of the Lower to Middle
1568 Paleolithic boundary: new dates from the Levant. Journal of Human Evolution 43, 107-
1569 122.

- 1570 Porat, N., Ronen, A., 2002. Luminescence and ESR Age Determinations of the Lower Paleolithic
1571 Site Evron Quarry, Israel. *Advances in ESR Applications* 18, 123-130.
- 1572 Potts, R., 1998. Environmental hypotheses of hominin evolution. *Yearbook of Physical*
1573 *Anthropology* 41, 93-136.
- 1574 Potts, R., 2002. Complexity and adaptability in human evolution. In: Goodman, M., Moffat, A.S.
1575 (Eds.), *Probing Human Origins*. American Academy of Arts and Sciences, Cambridge,
1576 MA, pp. 33-58.
- 1577 Rabinovich, R., Gaudzinski-Windheuser, S., Goren-Inbar, N., 2008. Systematic butchering of
1578 fallow deer (*Dama*) at the early middle Pleistocene Acheulian site of Gesher Benot
1579 Ya'aqov (Israel). *Journal of Human Evolution* 54, 134-149.
- 1580 Raia, P., Piras, P., Kotsakis, T., 2006. Detection of Plio-Quaternary large mammal communities
1581 of Italy. An integration of fossil faunas biochronology and similarity. *Quaternary Science*
1582 *Reviews* 25, 846-854.
- 1583 Rink, J.W., Bartoll, J., Schwarcz, H., Shane, P., Bar-Yosef, O., 2007. Testing the reliability of
1584 ESR dating of optically exposed buried quartz sediments. *Radiation Measurements* 42,
1585 1618-1626.
- 1586 Rightmire, G.P., Lordkipanidze, D., Vekua, A., 2006. Anatomical descriptions, comparative
1587 studies and evolutionary significance of the hominin skulls from Dmanisi, Republic of
1588 Georgia. *Journal of Human Evolution* 50, 115-141.
- 1589 Robinson, C., Belmaker, M., In Press. The biogeography of *Giraffa* spp. in the Southern Levant
1590 and its implication for 'Out of Africa I'. *PaleoAnthropology*.
- 1591 Rollefson, G.O., Schnurrenberger, D., Quintero, L.A., Watson, R.P., Low, R., 1997. 'Ain Soda
1592 and 'Ain Qasiya: New Late Pleistocene and Early Holocene sites in the Azraq Shishan

- 1593 area, Eastern Jordan. In: Gebel, H.G., Kafafi, Z., O., R.G. (Eds.), The Prehistory of
1594 Jordan II, Perspectives from 1996. ex oriente, Berlin, pp. 45-58.
- 1595 Ron, H., Gvirtzman, G., 2001. Magnetostratigraphy of Ruhama bad-land Quaternary deposit: a
1596 new age of the Lower Paleolithic site. (Eds.), Abstracts of the Annual Meeting of the
1597 Israel Geological Society. pp. 95.
- 1598 Ron, H., Levi, S., 2001. When did hominids first leave Africa? New high-resolution
1599 magnetostratigraphy from Erk-el-Ahmar Formation, Israel. *Geology* 29, 887-890.
- 1600 Ron, H., Porat, N., Ronen, A., Tchernov, E., Horwitz, L.K., 2003. Magnetostratigraphy of the
1601 Evron Member - Implication for the age of the Middle Acheulian site of the Evron
1602 Quarry. *Journal of Human Evolution* 44, 633-639.
- 1603 Ronen, A., 1991a. The Yiron gravel lithic assemblage, artefacts older than 2.4 Ma in Israel.
1604 *Archäologisches Korrespondenzblatt* 21, 159-164.
- 1605 Ronen, A., 1991b. The Lower Palaeolithic site Evron-Quarry in western Galilee, Israel.
1606 *Sonderveröffentlichungen Geologisches Institut der Universität zu Köln* 82, 187-212.
- 1607 Ronen, A., Inbar, M., Klein, M., Brunnacker, K., 1980. Artifact-bearing gravels beneath the
1608 Yir'on Basalt. *Israel Journal of Earth Sciences* 29, 221-226.
- 1609 Ronen, A., Burdukiewicz, J.-M., Laukhin, S.A., Winter, Y., Tsatskin, A., Dayan, T., Kulikov,
1610 O.A., Vlasov, V.K., Semenov, V.V., 1998. The Lower Palaeolithic Site Bizat Ruhama in
1611 the Northern Negev, Israel. Preliminary Report, 1996 Excavations. *Archäologisches*
1612 *Korrespondenzblatt RGZM* 28, 163-173.
- 1613 Rosenfeld, A., Nathan, Y., Feldman, H.R., Goldman, S. Yechieli, S.J. and Elkin, E. 2004.
1614 Paleoeology and paleosalinity of some Pleistocene lucustrine strata in Northern Israel:
1615 Acheulian culture, Trace elements and stable isotopes of ostracode valves. *Geological*

- 1616 Society of American Abstracts with Program Vol. 36 (2). P. 69.
- 1617 Rowland, J.C., 2006. Inter-and intra- tooth isotopic variation in mammalian tooth enamel from
1618 western Israel: Implication for paleoenvironmental and paleoclimate change over the past
1619 350 Kya. M.Sc. Dissertation, The University of Arizona.
- 1620 Sagi, A., 2005. Magnetostratigraphy of 'Ubeidiya Formation, Northern Dead Sea Transform,
1621 Israel. M.Sc. Dissertation, The Hebrew University.
- 1622 Sanlaville, P., 1988. Synthèse sur le paléoenvironnement. *Paléorient* 14, 57-60.
- 1623 Sanlaville, P., Besançon, J., Copeland, L., Muhesen, S., 1993. Le Paléolithique de la vallée
1624 moyenne de l'Oronte (Syrie). (Eds.), BAR International Series 587. BAR, Oxford.
- 1625 Sen, S., de Bonis, L., Dalfes, N., Geraads, D., Jaeger, J.J., Mazin, J.-M., 1991. Première
1626 découverte d'un site à mammifères pléistocènes dans une fissure karstique en Anatolie
1627 centrale. *C.R. Acad. Sci. Paris* 313, 127-132.
- 1628 Smith, P. E. L. (1986). *Paleolithic Archaeology in Iran*. American Institute of Iranian Studies.
1629 Museum University of Pennsylvania, Philadelphia.
- 1630 Sohn, S., Wolpoff, M.H., 1993. Zuttiyeh face: A view from the east. *American Journal of*
1631 *Physical Anthropology* 91, 325-347.
- 1632 Steudel, K.L., 1994. Locomotor energetics and hominid evolution. *Evolutionary Anthropology* 3,
1633 42-48.
- 1634 Stekelis, M., 1960. The Palaeolithic Deposits of Jisr Banat Yaqub. *The Bulletin of the Research*
1635 *Council of Israel* 9G, 61-90.
- 1636 Stiner, M. Clark Howell, F., Martinez-Navarro, B., Tchernov, E, and Bar-Yosef, O., 2001.
1637 Outside Africa: Middle Pleistocene Lycoan from Hayonim Cave, Israel. *Bulletino della*
1638 *Società Paleontologica Italiana* 40, 392-302.

1639 Stiner, M., R. Barkai, A. Gopher, 2004. The Acheulo-Yabrudian faunas from Qese Cave (Israel):
1640 Preliminary results. *PaleoAnthropolgy* 2, A52

1641 Stiner, M.C., Barkai, R., Gopher, A., 2009. Cooperative hunting and meat sharing 400-200 kya
1642 at Qesem Cave, Israel. *Proceedings of the National Academy of Sciences* 106, 13207-
1643 13212.

1644 Suc, J.-P., 1984. Origin and evolution of the Mediterranean vegetation and climate in Europe.
1645 *Nature* 307, 429-432.

1646 Spiro, B., Ashkenazi, S., Mienis, H.K., Melamed, Y., Fiebel, C. Delgado, A., Stratinsky, A.
1647 2009. Climate variability in the Upper Jordan Valley around 0.78 Ma, inference from
1648 time-series stable isotopes of Viviparidae, supported by mollusk and plant paleoecology.
1649 *Palaeogeography, Palaeoclimatology, Palaeoecology* 282: 32-44.

1650 Taskiran, H., 1998. The Distribution of Bifaces in Anatolia. In: Otte, M. (Ed.), *Préhistoire*
1651 *d'Anatolie: Genèse de deux mondes, Anatolian Prehistory: At the Crossroads of Two*
1652 *Worlds, Volume 2. Etudes et Recherches Archéologiques de l'Université de Liège, Liège,*
1653 *pp. 569-578.*

1654 Tappen, M., 2009. The wisdom of the aged and out of Africa I. In: J.J, S., Lieberman, D.E.
1655 (Eds.), *Transitions in Prehistory: Essays in Honor of Ofer Bar Yosef. American School of*
1656 *Prehistoric Research, pp. 24-41.85.*

1657 Tappen, M., Adler, D., Ferring, C.R., Gabunia, M., Vekua, A., Swisher, C.C., 2002. Akhalkalaki:
1658 The taphonomy of an early Pleistocene locality in the Republic of Georgia. *Journal of*
1659 *Archaeological Science* 29, 1367-1391.

1660 Tchernov, E., 1987. The age of the `Ubeidiya Formation, an early hominid site in the Jordan
1661 Valley. *Israel Journal of Earth Sciences* 36, 3-36.

- 1662 Tchernov, E., 1988. The biogeographical history of the southern Levant. In: Yom-Tov, Y.,
1663 Tchernov, E. (Eds.), The Zoogeography of Israel. Dr. Junk Publishers, Dordrecht, pp.
1664 159-250.
- 1665 Tchernov, E., 1994. New comments on the biostratigraphy of the Middle and Upper Pleistocene
1666 of the southern Levant. In: Bar-Yosef, O., Kra, R.S. (Eds.), Late Quaternary Chronology
1667 and Paleoclimates of the Eastern Mediterranean. pp. 333-350.
- 1668 Tchernov, E., 1995. The earliest hominids in the Southern Levant. In: Gibert, J., Sánchez, F.,
1669 Gibert, L., F., R. (Eds.), The Hominids and their Environments During the Lower and
1670 Middle Pleistocene of Eurasia. Ayuntamiento de Orce, Granada, pp. 389-406.
- 1671 Tchernov, E., Volokita, M., 1986. Insectivores and Primates from the early Pleistocene of
1672 'Ubeidiya Formation. In: Tchernov, E. (Ed.), Les Mammifères du Pléistocène Inférieur,
1673 de la Vallée du Jourdain à Oubéidiyeh. Association Paléorient, Paris, pp. 54-62.
- 1674 Tchernov, E., Kolska Horwitz, L., Ronen, A., Lister, A., 1994. The faunal remains from Evron
1675 Quarry in relation to other Paleolithic hominid sites in the Southern Levant. Quaternary
1676 Research 42, 328-339.
- 1677 Tchernov, E., Tsoukala, E., 1997. Middle Pleistocene (Early Toringian) carnivore remains from
1678 Northern Israel. Quaternary Research 48, 122-136.
- 1679 Tchernov, E., Belmaker, M., 2004. The biogeographic history of the fauna in the Southern
1680 Levant and the correlation with climatic change. In: Dolev, A., Perevolototsky, A. (Eds.),
1681 The Red Book. Vertebrates of Israel. The Israel Nature and Park Authority and the
1682 Society for the Protection of Nature in Israel, Jerusalem, pp. 21-25.
- 1683 Thomas, H., 1985. The Early and Middle Miocene land connection of the Afro-Arabian plateau
1684 and Asia: A major event for hominid dispersal? In: Delson, E. (Ed.), Ancestors: The Hard

- 1685 Evidence. A. R. Liss, New York, pp. 42-50.
- 1686 Thomas, H., Geraads, D., Janjou, D., Vaslet, D., Memesh, A., Billiou, D., Boucherns, H.,
 1687 Dobigny, G., Eisenmann, V., Gayet, M., Lapparent De Broin, F., Petter, G., Halawani,
 1688 M., 1998. First Pleistocene faunas from the Arabian Peninsula: An Nafud desert, Saudi
 1689 Arabia. *Comptes rendue de l'Academie des science, série II de la terre et des planètes*
 1690 326, 145-152.
- 1691 Tobias, P., 1966a. A Member of the Genus *Homo* from 'Ubeidiya. *Publications of the Israel*
 1692 *Academy of Sciences and Humanities*, Jerusalem.
- 1693 Tobias, P.V., 1966b. Fossil hominid remains from 'Ubeidiya, Israel. *Nature* 211, 130-133.
- 1694 Tzedakis, P.C., 2007. Seven ambiguities on the Mediterranean paleoenvironmental narrative.
 1695 *Quaternary Science Reviews* 26, 2042-2066.
- 1696 Tzedakis, P.C., Hooghiemstra, H., Pälike, H., 2006. The last 1.35 million years at Tenaghi
 1697 Philippon: revised chronostratigraphy and long-term vegetation trends. *Quaternary*
 1698 *Science Reviews* 25, 3416-3430.
- 1699 Vaks, A., Bar-Matthews, M., Ayalon, A., Frumkin, A., 2002. Paleoclimate Conditions in the
 1700 Central and Northern Negev Desert During the Pleistocene. 2002 Ma'agan Israeli
 1701 *Geological Society Abstracts* 125.
- 1702 van Zeist, W., Bottema, S., 2009. A palynological study of the Acheulian site of Gesher Benot
 1703 Ya'aqov, Israel. *Vegetation History and Archaeobotany* 18, 105-121.
- 1704 Vandermeersch, B., 1995. Le rôle du Levant dans l'évolution de l'humanité au Pléistocene
 1705 supérieur. *Paléorient* 21, 25-34.
- 1706 Vekua, A., 1986. Pleistocene Mammalian Fauna of Akhalkalaki (Southern Georgia, USSR).
 1707 *Paleontographia Italica* 74, 63-96.

- 1708 Vekua, A.K., 1987. Lower Pleistocene Mammalian fauna of Akhalkalaki (Southern Georgia,
1709 USSR). *Palaeontographica Italica* 74, 63-96.
- 1710 Vekua, A., 1995. Die Wirbeltierfauna des Villafranchium von Dmanisi und ihr
1711 biostratigraphische Bedeutung. *Jahrbuch des Roemisch-Germanischen Zentralmuseums*,
1712 Mainz.
- 1713 Vekua, A., Lordkipandize, D., 2008. The history of vertebrate fauna in Eastern Georgia. *Bulletin*
1714 *of the Georgian National Academy of Sciences* 2, 149-155.
- 1715 Verosub, K., Tchernov, E., 1991. Resultats préliminaires de l'étude magnétostratigraphique d'une
1716 séquence sédimentaire à l'industrie humaine en Israël. In: Vandermeersch, B. (Ed.), *Les*
1717 *premiers peuplements de l'Europe*. C.N.R.S, Paris, pp. 237-242.
- 1718 von Koenigswald, W., 1998. Preliminary report on the middle Pleistocene small mammal fauna
1719 from Yarimburgaz cave in Turkish Thrace. *Eiszeitalter und Gegenwart* 48, 162-169.
- 1720 Vrba, E.S., 1988. Late Pliocene climatic events and hominid evolution. In: Grine, F.E. (Ed.),
1721 *Evolutionary History of the "Robust" Australopithecines*. Aldine de Gruyter, New York,
1722 pp. 405-426.
- 1723 Vrba, E.S., 1995. The fossil record of African antelopes (Mammalia, Bovidae) in relation to
1724 human evolution and paleoclimate. In: Vrba, E., Denton, G., Partridge, T., Burckle, L.
1725 (Eds.), *Paleoclimate and Evolution, with Emphasis on Human Origins*. Yale University
1726 Press, New Haven, pp. 385-424.
- 1727 Walker, A., Leakey, R., (Eds.) 1993. *The Nariokotome Homo erectus Skeleton*. Harvard
1728 University Press, Cambridge.
- 1729 Walker, A., Shipman, P., 1996. *The Wisdom of Bones: In Search of Human Origins*. Knopf,
1730 New York.

- 1731 Weinstein-Evron, M., Bar-Oz, G., Zaidner, Y., Tsatskin, A., Druck, D., Porat, N., HersHKovitz,
1732 I., 2003. Introducing Misliya Cave, Mount Carmel, Israel: a new continuous
1733 Lower/Middle Paleolithic sequence in the Levant. *Eurasian Prehistory* 1, 31-55.
- 1734 Werker, E. (2006). 780,000-year old wood from Gesher Benot Ya'akov, Israel. *Israel Journal of*
1735 *Plant Sciences* 54, 291-300.
- 1736 Whalen, N., Sindi, H., Wahida, G., Siraj-Ali, J.S., 1983. Excavations of Acheulean sites near
1737 Saffaqah in Ad-Dawadmi 1302-1982. *Atlal* 7, 9-21.
- 1738 Whalen, N., Siraj-Ali, J.S., Davis, W., 1984. Excavations of Acheulean sites near Saffaqah,
1739 Saudi Arabia, 1403 AH 1983. *Atlal* 8, 9-24.
- 1740 Wynn, J.G., 2004. Influence of Plio-Pleistocene aridification on human evolution: Evidence from
1741 paleosols of the Turkana Basin, Kenya. *American Journal of Physical Anthropology* 123,
1742 106-118.
- 1743 Zaidner, Y., 2003a. The use of raw material at the Lower Palaeolithic site of Bizat Ruhama,
1744 Israel. In: Burdukiewicz, J.M., Ronen, A. (Eds.), *Lower Palaeolithic Small Tools in*
1745 *Europe and the Levant*. BAR, Oxford, pp. 121-131.
- 1746 Zaidner, Y., 2003b. The lithic assemblage of Bizat Ruhama: Lower Paleolithic site in southern
1747 coastal plain, Israel. M.Sc. Dissertation, Haifa University, Israel.
- 1748 Zaidner, Y., Ronen, A., Burdukiewicz, J.,-M., 2003. L'industrie microlithique de Paléolithique
1749 inférieur de Bizat Ruhama, Israël. *L'Anthropologie* 107, 203-222.
- 1750 Zarins, J., Ibrahim, M., Potts, D., Edens, C., 1979. Saudi Arabian archaeological reconnaissance
1751 1978: The preliminary report on the third phase of the comprehensive archaeological
1752 survey program - the central province. *Atlal* 3, 9-38.
- 1753 Zarins, J., Whalen, N., Ibrahim, M., al Jawad, A., Khan, M., Khan, M., 1980. Saudi Arabian

1754 archaeological reconnaissance 1979: Preliminary report on the central and southwest
 1755 provinces survey. Atlal 4, 9-36.

1756 Zarins, J., Rahbini, A., Kamal, M., 1982. Preliminary report on the archaeological survey of the
 1757 Riyadh area. Atlal 6, 25-38.

1758 Zazanashvili, N., Gagnidze, R., Nakhutsrishvili, G., 2000. Main types of vegetation zonation on
 1759 the mountains of the Caucasus. Proceedings IAVS symposium pp. 214-217.

1760 Zhu, R.X., An, Z., Potts, R., Hoffman, K.A., 2003. Manganetostratigraphic dating of early
 1761 humans in China. Earth Science Reviews 61, 341-359.

1762 Zohary, D., 1973. Geobotanical foundations of the Middle East. Gustav Fisher,
 1763
 1764
 1765

1765 **Figure legends**


1766

1767 Figure 1: Location of major sites mentioned in the texts. Subfigure A includes the near east.


1768 Subfigure B includes the Levant and is denoted in the box within subfigure A.

1769

1770



A.



	<i>Panthera leo</i>								X		
	<i>Panthera gombaszoegensis</i>	X		X							

	<i>Giraffa</i> sp.		X								
	<i>Giraffa camelopardalis</i>				X						
Bovidae	<i>Bison georgicus</i>	X									
	<i>Leptobos</i> sp.		X								
	<i>Bison</i> sp.				X		X		X	X	
	<i>Bos primigenius</i>				X		X		X	X	X
	<i>Gallogoral meneghini</i>	X									
	<i>Pelorovis oldowayensis</i>		X								
	<i>Bos buiaensis</i>							X			
	? <i>Sinoreas</i> sp.						X				
	<i>Capra</i> sp.							X			
	<i>Capra ibex</i>								X		X
	<i>Capra aegagrus</i>										X
	<i>Capra dalii</i>	X									
	<i>Caprini</i> indet.							X			
	<i>Ovibovini</i> indet.	X						X			
	<i>Pontoceros</i> sp.	X	X		X						
	<i>Antilopini</i> indet.	X	X								
	<i>Gazella</i> sp.		X		X			X			
	<i>Gazella</i> cf. <i>gazella</i>								X		X
	<i>Oryx</i> cf. <i>gazella</i>		X								
	<i>Alcelaphus buselaphus</i>				X						X
Camelidae	<i>Camelus</i> sp.	X	X		X						
Suidae	<i>Kolpochoerus olduvaiensis</i>		X								
	<i>Kolpochoerus evronensis</i>				X						
	<i>Sus strozzi</i>		X								
	<i>Sus scrofa</i>							X	X		X
Hippopotamidae	<i>Hippopotamus georgicus</i>	X?					X				
	<i>Hippopotamus behemoth</i>		X		X						
	<i>Hippopotamus gorgops</i>		X								
	<i>Hippopotamus antiquus</i>				X			X			X
Primates	<i>Macaca sylvanus</i>		X								
	Cercopithecidae cf. <i>Theropithecus</i>		X								

Table 2: Faunal and hominin dispersal from Africa to Eurasia in different biozones

Date (Ma)	MOIS	Mammalian zone	Caucasus	Anatolia	Levant	African Dispersal ¹	Hominin dispersals	Cultural entities
2.5		Early Villafranchian	Kvebebi		Bethlehem	N=2		
1.8		Middle Villafranchian	Dmanisi, Tsalka				Dispersal event	Core and flake
1.6 – 1.2		Late Villafranchian			‘Ubeidiya	N=5	Dispersal event	Early Acheulian
1.1 – 1.0		Epi-villafranchian	Akhalkalaki Takshe	Dursunlu	Daquara, Latamne, Evron,	N=2		
0.8	19	Gallerian			GBY	N=1	Dispersal event	Acheulian
0.5				Danizli, Emirkaya -2, Bear Cave,	Revadim Holon, Ein Soda, Tel Hesi			
0.4 – 0.3	8-9	Auralian			Qesem, Revadim Tabun E			Achelo-Yaburidian
0.3 – 0.13	7-6				Hayonim E, Misliya			Early Mousterian

¹ Number of species that left Africa

